



British  
Orthopaedic  
Association



# **Best Practice for Management of Anterior Cruciate Ligament (ACL) Injuries**

**Review date- Sep 2023**

**BOA, BASK, BOSTAA Elective Care Standards:****Best Practice for Management of Anterior Cruciate Ligament (ACL) Injuries**

British Association for Surgery of the Knee (BASK) and British Orthopaedic Sports Trauma and Arthroscopy Association (BOSTAA) have collaborated to commission a committee to produce best practice guidelines on the management of Anterior Cruciate Ligament (ACL) injuries. The Guideline Development Group (GDG) is made up as follows:

<b>Mr Sanjeev Anand</b>	<b>Chairperson, Editor</b>	<b>Consultant Orthopaedic Surgeon, Leeds Teaching Hospitals NHS Trust</b>
<b>Mr Panos Thomas</b>	<b>Chairperson, Editor</b>	<b>Consultant Orthopaedic Surgeon, Whittington Hospital NHS Trust, London</b>
Mr Simon Roberts	Outpatient Consultation	Consultant Orthopaedic Surgeon, Robert Jones & Agnes Hunt Hospital, Oswestry
Dr Philip Robinson	Imaging	Consultant Musculoskeletal radiologist, Leeds Teaching Hospitals NHS Trust;
Mr Mark Blyth	Indications for the operation	Consultant Orthopaedic Surgeon, Glasgow Royal Infirmary
Mr Andrew Porteous	Management of associated peripheral injuries	Consultant Orthopaedic Surgeon, North Bristol NHS Trust
Mr James Robinson		Consultant Orthopaedic Surgeon, North Bristol NHS Trust
Mr Paul Gibb	Paediatric Injuries	Consultant Orthopaedic Surgeon, Maidstone and Tunbridge Wells NHS Trust
Mr Mike Carmont	Preoperative assessment	Consultant Orthopaedic Surgeon, Shrewsbury and Telford NHS Trust
Prof Deiry Kader	Admission and thromboprophylaxis	Consultant Orthopaedic Surgeon, South West Elective Orthopaedic Centre, Epsom
Mr Tony Hui	Hospital & theatre resources	Consultant Orthopaedic Surgeon, Middlesbrough
Dr Chris Hargreaves	Anaesthesia	Consultant Orthopaedic Surgeon, Whittington Hospital NHS Trust

Mr Richard Parkinson	The Surgeon	Consultant Orthopaedic Surgeon, Wirral University Teaching Hospital
Prof Fares Haddad	Surgical Procedure	Consultant Orthopaedic Surgeon, University College Hospitals, London
Mr Adil Ajuied	Postoperative management	Consultant Orthopaedic Surgeon, Guy's and St Thomas' NHS Foundation Trust, London
Mr Sean O'Leary	Ligament Registry	Consultant Orthopaedic Surgeon, Royal Berkshire, Hospital, Reading
Dr Lee Herrington	Conservative management and Rehabilitation	Physiotherapist & Lecturer in Sports Injury rehabilitation, Salford

**With Editorial input from: Prof Philip Turner, Immediate Past President, BOA**

**Prof Andrew Price, President, BASK**

**Prof Fares Haddad, President, BOSTAA**

## **DISCLAIMER**

The recommendations in this guideline represent the view of the GDG, arrived at after careful consideration of the evidence available. It represents a consensus statement from The British Association for Surgery of the Knee and the British Orthopaedic Sports Trauma and Arthroscopy Association. It is hoped that this Guide will inform surgeons, hospitals and commissioners in making decisions about the care and management of patients. When exercising their judgement, professionals are expected to take this guideline fully into account, alongside the individual needs, preferences and values of their patients or service users. The application of the recommendations in this guideline is not mandatory and the guideline does not override the responsibility of healthcare professionals to make decisions appropriate to the circumstances of the individual patient, in consultation with the patient and/or their carer or guardian.

## **FOREWORD**

We are delighted to support the dissemination of this document on best practice for management of anterior cruciate ligament injuries. These are extremely important injuries that affect more than 20,000 active individuals in the UK every year. Timely and appropriate management is critical to allow patients to regain their quality of life and activity and to try and help them to preserve function. The clinical guidance in this document spans the entire patient pathway and will act to improve care at all stages.

This document represents a collaboration between the British Association for Surgery to the Knee and the British Orthopaedic Sports Trauma and Arthroscopy Association with renowned experts summarising the current literature and current thinking and highlighting optimal management strategies. The work had been supported by the British Orthopaedic Association and is an important contribution to the growing body of best practice guidance developed with their help. We hope this will be one of several ongoing collaborations between the respective specialist societies and that it will benefit surgeons, allied practitioners and most importantly our patients.

Fares Haddad, President of BOSTAA

Andrew Price, President of BASK

## Table of Contents

## Page

SUMMARY OF AUDIT STANDARDS .....	1
1. Introduction .....	3
2. The outpatient consultation.....	3
3. Radiology.....	5
4. Indications for operations.....	7
5. Management of associated peripheral and other injuries.....	9
6. Juvenile ACL.....	13
7. Preoperative Assessment.....	17
8. Admission to hospital.....	19
9. Hospital facilities required.....	19
10. Anaesthesia.....	20
11. Surgeon.....	23
12. Thromboprophylaxis.....	24
13. Surgical Procedure.....	25
14. Postoperative care and followup.....	28
15. National Ligament Registry.....	29
16. Conservative management, rehabilitation and return to sport.....	31
17. Bibliography.....	34

**BOA, BASK, BOSTAA Elective Care Standards: Best Practice for Management of Anterior Cruciate Ligament (ACL) Injuries**

**Summary of Audit Standards**

1. Patients with haemarthrosis, following an acute knee injury, should be assessed by a clinician proficient in assessment of knee injuries. This should be in form of an Acute Knee injury clinic led either by a surgeon, a physiotherapist or a suitably trained doctor or allied professional.

a. This assessment should be done within 2 weeks of presentation, so as to identify injuries requiring urgent treatment.

b. Urgent access to imaging facilities should be available from this clinic. Magnetic resonance imaging (MRI), reported by a musculoskeletal radiologist, remains the imaging modality of choice to assess for ACL injury and associated internal derangement.

c. All coexisting injuries should be identified and documented, following imaging and repeat assessment, as required.

d. A management plan should be made in consultation with the patient after discussion of the treatment options, both non operative and operative.

2. The prime indication for ACL reconstruction is symptomatic instability. The decision for early surgical reconstruction vis a vis trial of non-operative treatment should be individual to each patient.

3. All patients being considered for surgery should be offered prehabilitation to recover knee movement and quadriceps strength.

4. Surgery, if considered, is usually performed when the knee is “quiet” with no swelling and a full range of movement, particularly extension, has been restored.

5. Patients with an unstable and repairable meniscal lesion in combination with ACL insufficiency should be ideally offered early combined meniscal repair and ACL reconstruction. Staged meniscal repair followed by later ACL reconstruction is acceptable in the case where the patient presents acutely with a locked knee associated with an ACL rupture and to allow the reconstruction to proceed when the knee has regained full extension or when suitable surgical expertise is available.

6. ACL injuries in patients who have not reached skeletal maturity and multiligament injuries should be managed by surgeons with an interest in the field.

**7. Consent**

a. Non-operative and operative options should be discussed. The benefits, limitations and likely outcome of each should be discussed and recorded. The general risks of surgical intervention together with the specific risks of reconstructive surgery should be discussed and documented. Risks of persistent instability and reinjury, including to the opposite knee should be discussed. Patients should be advised on timing of return to sports, which should be specific to the individual, but should rarely be less than 9 months after surgery due to higher risk of reinjury.

b. Graft selection should be discussed including autograft, allograft and synthetic ligaments. This should include the benefits, complications and risks of all and the preferred graft

recommendation.

c. The management of chondral and meniscal pathology should be discussed together with the implications on post-operative recovery.

d. Consent for inclusion into National Ligament Registry should be sought and patients should be registered in its database. Hospitals should facilitate the accurate recording of surgical procedures and patients' outcome by providing appropriate clerical and IT support.

## **8. Surgery**

a. The procedure should be performed on a Day Case basis, for majority of patients. The surgery should be performed by or under supervision of a surgeon with special interest in soft tissue knee reconstruction.

b. Local anaesthetic infiltration, adductor canal block or femoral nerve block may be used for perioperative pain relief. When hamstring graft is used, harvest site should be infiltrated with local anaesthetic.

c. Pharmacological VTE prophylaxis is not indicated as a routine. However, all patients should be risk assessed upon admission to hospital. In high risk cases, chemical thromboprophylaxis may be used as per local guidelines. Mechanical prophylaxis like calf pumps should be used intraoperatively and in recovery following surgery, before the patient becomes mobile.

d. An examination under anaesthetic must be performed to take into account the degree of anteroposterior and rotational laxity as well as any other associated injuries and documented.

e. Graft choice should be decided based on individual patient characteristics and surgeon experience. Allografts are not recommended for primary reconstructions in younger patients (<35 yrs old). Synthetic ligaments are not currently recommended for routine primary reconstruction.

f. Tunnel position is a source of debate but any reconstruction should allow full range of movement of the knee joint with no impingement in the notch. There should be no excessive motion of graft on knee movement. Stability should be restored and pivot should be abolished following reconstruction. Graft fixation technique and implants should allow immediate knee mobilisation.

9. Following surgery, early quadriceps activation and full range motion should be encouraged with support from an appropriate physiotherapy programme.

10. Decision to return to sport should be criteria based taking into consideration physical factors relating to the knee; psychological factors including fear of reinjury and social factors; while being tailored to the specific sport. To assess readiness to return to play and the risk for reinjury, a range of tests, including strength tests, hop tests and measurement of movement quality, should be used.



## **1 INTRODUCTION**

- 1.1 Anterior cruciate ligament (ACL) injury is a common injury usually affecting young and active individuals. It is the commonest knee injury requiring surgical intervention(1). Evidence on management of this injury is evolving. Though good outcome is achieved in majority of patients, still a large proportion of patients are unable to return to their preinjury state(2). The final outcome for patients is dictated not just by injury to the ACL but also by associated injuries to the knee joint. This is further influenced by a variety of psychosocial factors, including patient's aspirations and demands. Clinicians looking after these patients have an opportunity to shape the outcome by influencing the whole package of care; from the assessment at presentation, to treatment (both surgical and non-surgical) and rehabilitation, leading to the desired functional outcome for the patient.
- 1.2 This guide collates the current evidence on management of ACL injuries and summarises treatment options for the benefit of involved clinicians. These guidelines have been produced in collaboration between the BASK and BOSTAA. It may not be applicable to all patients and in all circumstances. The treating clinicians should consider the individual requirements of each patient, before formulating a management plan.

## **2. THE OUTPATIENT CONSULTATION**

- 2.1 The orthopaedic outpatient consultation allows exchange of information between the patient and a clinician who has experience and expertise in both conservative & surgical management of patients with ACL injury to inform the shared decision-making process. There should be liaison between primary and secondary care in planning the local delivery of acute knee services, but it is considered best practice for all post-traumatic knee haemarthrosis to be assessed urgently by a clinician with special expertise in knee injury management(3–5).
- 2.2 Whether the ACL is deficient, what other comorbidity exists, and the extent to which that deficiency is responsible for current symptoms is determined by an assessment of the history and physical examination, usually supplemented by imaging.

- 2.3 It should take place face to face in a confidential environment, with access for relatives and allied health professionals with a minimum time of 15-20 minutes. Patients will often be months or even years after injury and frustrated by the delay in diagnosis. It may be the first time the diagnosis is made, but equally they may already have researched the diagnosis and made a decision on their preferred treatment. This decision may change following the consultation.
- 2.4 Past medical records from hospital and general practitioner including imaging (if any) should be available.
- 2.5 The history of the injury, subsequent and current symptoms should be documented, followed by a clinical examination and review of all imaging. Instrumented laxiometry may be helpful but is not considered to be universally necessary.
- 2.6 A management plan is made after discussion of both operative & non-operative options for both the ACL and coexisting knee pathologies, including the role of rehabilitation, whether pre or post-operative or as definitive treatment, bracing, modification of activities and surgical repair or reconstruction(6–13).
- 2.7 This is the start of the process of informed consent. The intended and likely benefits, along with serious or frequently occurring risks of harm or failure of treatment are discussed at a level suitable to the patient. This should include the likelihood of stability and a successful return to sporting activities being achieved in the short term, as well as the long term effect of instability on the knee, its menisci & articular surface(2,11,14–17). Patients should be aware of the lack of evidence for reconstruction reducing the risk osteoarthritis or improving long-term function in the absence of instability(15,18–21).
- 2.8 The arrangements for treatment, postoperative management & follow up are described with an estimate of duration, along with the role of the National Ligament Registry need to be discussed, but this may best be delegated to an allied health professional & supplemented with written and internet documents.

### 3. RADIOLOGY

- 3.1 Although all patients will probably have plain x-rays, Magnetic resonance Imaging (MRI) remains the imaging modality of choice to assess the knee for ACL injury and associated internal derangement.
- 3.2 Ideally the MRI should be reported by an experienced musculoskeletal radiologist who works with the referring clinical team.
- 3.3 MRI can now be 1.5T and 3.0T magnet strength, with lower strength magnets (<1.0T) less frequently used and unproven for accuracy. In the UK most clinical scanners are currently 1.5T with the incidence of 3.0T increasing.
- 3.4 Patients should lie supine with the knee minimally flexed encased in a specific multi-channel knee coil (typically at least 8 channel).
- 3.5 Typically, the MRI protocol can be comprehensive and still total less than 25 minutes allowing quick throughput. Longer scanning times with complex protocols often result in more patient discomfort and increased movement artefacts.
- 3.6 MRI Protocols should include(22–24);
  - 3.6.1 Anatomical planes to include; sagittal oblique (angled 10 degrees so parallel to the ACL), coronal and transverse axial. Additional ACL specific coronal oblique sequences have been proposed but are rarely used in clinical practice and are not of proven diagnostic benefit. Specific orientations for the posterolateral corner have also been suggested but again rarely produce significant additional information to the standard planes and sequences. Newer faster 3D volume sequences (6-8 minutes) allow reformatting in any anatomical plane with good resolution particularly on 3.0T scanners.
  - 3.6.2 Pulse sequences to include; PD weighted (+/- fat suppression), T2 weighted fat (+/- fat suppression) and T1 weighted (not fat suppressed) sequences to optimally assess all

structures including bone, articular cartilage, menisci and ligaments. Newer faster 3D volume sequences have been assessed and show promising results compared to traditional sequences for diagnostic accuracy in assessing menisci and ligaments particularly on 3.0T scanners but have not replaced established pulse sequences.

3.7 The origin, distal footprint and double bundle structure of the ACL is appreciated on MRI particularly on coronal and transverse axial sequences(22).

3.8 MRI has been shown to be very accurate for full thickness ACL tear (sensitivity >83-95% and specificity 95-100%). MRI findings can be classified as primary and secondary signs of ACL injury with primary signs the most specific(22–24).

3.8.1 Primary ACL injury signs include; ligament defect, diffuse abnormal signal, altered orientation or bowing (including collapse and pseudo mass) and non-visualisation.

3.8.2 Secondary signs include pivot shift pattern of osseous injuries, Segond fracture, anterior tibial translation (>5mm) and uncovering of the posterior horn of the lateral meniscus (>3.5mm).

3.9 MRI is accurate for assessing associated or combination injuries including bone bruising, Segond fracture, osteochondral lesions, menisci (tear or meniscocapsular injury), capsule, other ligament injuries and the posterolateral corner for oedema. Meniscal accuracy is the most studied and highly sensitive and specific for tears (>92%).

3.10 Bone marrow injury patterns can help determine the mechanism of injury, for example; pivot shift, hyperextension or varus rotation.

3.11 MRI accuracy is reduced in partial and chronic tears, as some primary signs described for complete tears are not present, such as orientation abnormality(22,23). Partial tear accuracy is reported at sensitivity >62-95% and specificity 19-97% but a lot of these studies are historic and used older equipment and sequences, hence the wide range reported. Recent studies on 3.0T scanners report sensitivity 77% and specificity 97%(25,26). Partial and chronic tears are best appreciated on coronal and transverse axial sequences(25,27).

3.12 Post reconstruction surgery - Caution is required within 18 months of surgery as the graft normally shows temporal signal change as it undergoes ligamentisation (synovial ingrowth and revascularisation); <3 months low T1w and T2w signal, 3-12 months increased T1w and T2w signal, then returning to low T1w and T2w signal by >18months(22,28). However the type of graft can also affect appearances with hamstring bundle reconstructions liable to have intervening fluid compared to patella grafts therefore resulting in a normal increase of T2w signal(28).

3.13 Post-surgical repair MRI assessment can evaluate for degenerative change, progressive osteochondral damage, graft failure, impingement lesions, tunnel abnormalities (cysts, expansion, lysis), extruded hardware, arthrofibrosis, donor site complications, acute graft tear or other acute internal derangement. Graft laxity while anatomically intact is very difficult to determine on imaging alone(23).

3.14 Artefact reducing sequences are now available to allow better assessment of tunnels and associated hardware on MRI. CT can be used when accurate tunnel sizes are required prior to revision.

3.15 MRI remains the imaging modality of choice to assess the knee for ACL injury and associated internal derangement. Accuracy is reduced in studies evaluating partial or chronic injuries but this often within a research setting. This can be minimised in clinical practice by correlation with the clinical history and examination findings from the referrer.

#### **4. THE INDICATIONS FOR THE OPERATION**

4.1 At this time, results of only two small randomised trials comparing early reconstruction to structured rehabilitation are available (29–31). While these studies suggest equal outcome between early reconstruction group, rehabilitation group and delayed reconstruction group; concerns remain over high cross over rates from non-operative arm to the surgical arm (9,29,31,32). There is also some concern about poorer long term outcome in the group undergoing delayed ACL reconstruction whilst reconstructed ACL appears to offer greater

objectively measured stability to the knee joint (13,29,32). ACL reconstruction also appears to have some protective effect on further meniscal injury, which may influence future osteoarthritis rates (11,33). Results from an ongoing UK based pragmatic randomized controlled trial are awaited (34).

4.2 Based on the currently available evidence, consensus group believes that decision for early surgical reconstruction vis a vis trial of non-operative treatment should be individual to the patient. This should be part of shared decision making with the patient, following discussion of their activity levels, aspirations and full assessment of the injury complex, including of associated injuries.

4.3. The aim of ACL reconstruction is to restore functional stability of the knee without compromising other joint functions, particularly range of movement(8). This is particularly important in the context of the knee with multiple ligament injury.

4.4 The prime indication for ACL reconstruction is symptomatic instability.

4.3 Each patient should be individually assessed taking into account, the type and frequency of physical activity and laxity at presentation, including a positive pivot shift test. Based on this assessment and following discussion with the patient, in certain circumstances primary reconstruction, before instability symptoms have been allowed to develop may be considered appropriate. This is usually once the initial inflammatory response to injury has settled, the knee is “quiet” with no swelling and a full range of movement, particularly extension, has been restored(8).

4.4 The presence of an unstable and repairable meniscal lesion in combination with ACL insufficiency is a strong indication for early combined ACL reconstruction and meniscal repair rather than partial meniscus excision. Staged meniscal repair followed by later ACL reconstruction is acceptable in the case where the patient presents with a locked knee associated with an ACL rupture, to allow the reconstruction to proceed when the knee has regained full extension(35).

4.5 There is no evidence as yet that reconstruction of the ACL reduces the incidence or progression of degenerative change in the knee, but early stabilization reduces the incidence

of subsequent meniscal pathology(11,33). Although no long term comparative studies are available, it is reasonable to conclude that this will have a protective effect(19).

4.6 Age and degenerative change are not in themselves contraindications to ACL reconstruction(36,37).

4.7 ACL injuries in patients who have not reached skeletal maturity pose particular problems, particularly in the very young, and should be managed by surgeons with an interest in the field.

## **5. MANAGEMENT OF ASSOCIATED PERIPHERAL AND OTHER INJURIES**

5.1 Injuries associated with Anterior Cruciate Ligament rupture include: injury to other knee ligaments, medial and lateral meniscal tears, and articular cartilage injuries. In a cohort study of 1145 consecutive patients with traumatic knee haemarthrosis who underwent MRI within 8 days after the injury, 52% had ACL injury (38). However only in 12% of cases, the ACL injury happened in isolation. 39% of patients had associated MCL injury, 10% LCL injury, 3% PCL injury, 55% meniscal tear, 31% cortical depression and 1% osteochondral fracture(38). The most serious of these is a multi-ligament injury or dislocation, which can be a limb threatening injury. Increased failure rate of ACL reconstruction surgery may occur in patients with missed posterolateral corner (39) or MCL injury(40,41). Associated meniscal and chondral injuries increase risk of poorer outcome, with higher risk of development of osteoarthritis(42,43). Delay in managing a displaced meniscal tear reduces the chance of the tear being successfully repaired.

5.2 Concomitant injury should be excluded by careful examination of all ligamentous structures and additional imaging that should include MRI scan (44–46) and, in acute injury, x-ray. Identification of associated injuries may significantly alter the urgency of intervention required in the ACL injured knee.

5.3 **Dislocation/Multiple ligament injury:** These are defined as complete rupture of 2 or more of the main knee ligaments, usually ACL and PCL They are usually the result of high-energy

vehicular trauma but can occur in high impact sports and in low-velocity injuries in hypermobile or obese patients (47–50).

- 5.3.1 Emergency reduction of true dislocations is required.
- 5.3.2 Assessment and documentation of pre and post reduction vascular status is required as vascular injury occurs in up to 50% of dislocations or bicruciate injuries (51,52). There is a higher risk of vascular injuries not just in high velocity injuries, but also in ultralow velocity knee dislocations (53,54).
- 5.3.3 Vascular surgeons should be involved urgently if limb perfusion is compromised.
- 5.3.4 Facilities for urgent vascular radiology investigation are required, or rapid referral to an appropriate unit considered.
- 5.3.5 If distal pulses are asymmetrical or and ankle brachial pressure index (ABPI) <0.9 vascular investigation (angiogram, CT angiogram, MR angiogram) should be performed. If distal pulses are present and ABPI >0.9 the patient should be admitted, serially examined for 24 hours and vascular investigation should be considered prior to ligament reconstruction(55).
- 5.3.6 In low-velocity injury, a well-perfused foot may be observed clinically but this does not necessarily exclude an intimal tear and any variation in vascular observations or ABPI <0.9 should prompt further investigation(55).
- 5.3.7 Spanning external fixator treatment may be required as part of emergency management in the setting of vascular injury/repair or inability to hold the joint reduced by other means(56).
- 5.3.8 Early discussion of these cases and referral/transfer to an appropriately specialized knee surgeon is necessary to allow planning of further investigation or intervention.

#### **5.4 Posteromedial Corner (PMC) Injury**

5.4.1 Injury to the PMC - superficial MCL (sMCL), the deep MCL (dMCL) and the postero-oblique ligament (POL) – should be assessed by valgus stress testing at 0° and 20° - 30° knee flexion(57). Stability to valgus stress in full extension indicates no significant damage to the posteromedial capsule / POL(58,59).

5.4.2 PMC injuries may be graded as:

Grade 1: localized medial pain with no appreciable valgus laxity.



Grade II: pain along the sMCL ligament with valgus laxity at 20° - 30° knee flexion, but not in extension,

Grade III: complete tear involving all three functional structure (sMCL, dMCL and POL) with valgus laxity in extension.

5.4.3 Failure to appreciate the rotational instability resulting from a torn PMC may lead to failure of ACL reconstruction(40,41,60).

5.4.4 Combined ACL rupture and MCL injury may be treated initially with functional bracing(57,61).

5.4.5 The brace can usually be discontinued 8 to 12 weeks after injury(57,61).

5.4.6 Delayed, isolated ACL reconstruction in patients with chronic ACL rupture and Grade 2 MCL laxity has shown similar postoperative ACL stability and outcome to isolated ACL injuries(62).

5.4.7 Earlier intervention for medial injury may be warranted in specific indications: severe Grade 3 MCL injuries, intra-articular entrapment of the ruptured MCL, large bony avulsion, complete tibial avulsion, the presence of antero-medial rotatory instability and patients who have valgus knee alignment(58,63,64).

## **5.5 Posterolateral corner (PLC) injury**

5.5.1 Posterolateral corner injury may be present in 9-10% of patients presenting with a knee haemarthrosis (38,65) and 10-15% of patients with chronic ACL deficiency(66).

5.5.2 Posterolateral laxity increases ACL graft failure(66–69).

5.5.3 Examination should include assessment of: limb alignment, gait, varus laxity at 0° and 30° flexion, and external rotation laxity at 30° and 90° (“Dial Test”)(70).

5.5.4 Grade 1 and 2 sprains may respond well to non-operative treatment; although residual laxity may remain. Conservatively managed grade 3 injuries are associated with high levels of osteoarthritis(63).

5.5.5 Repair or reconstruction of the PLC should be considered in the setting of acute or chronic posterolateral instability(71). Some reports suggest higher revision rates with primary repair (72,73) while more recent report suggest equally good outcome of repair when compared with reconstruction (74). Repair, if considered, should be performed within 2-3 weeks of the original injury (70)

5.5.6 High Tibial Osteotomy should be considered for chronic combined grade 3 posterolateral knee injuries and varus alignment(65,70).

## **5.6 Posterior Cruciate Ligament (PCL) Injury**

- 5.6.1 Combined ACL and PCL injury is uncommon but there should be a high index of suspicion that a dislocation has occurred and appropriate neurological and vascular assessment should be performed.
- 5.6.2 Superior results have been reported for surgical treatment of combined ACL / PCL injuries compared with conservative treatment(75–78). Evidence for superiority of a particular: surgical timing, technique or graft, is less clear.

## **5.7 Meniscal Injury**

- 5.7.1 Meniscal injury is common after ACL tear, with reported incidence varying between 30 and 80%. Acutely, lateral meniscal injury is more common, but medial meniscal pathology increases after ACL injury. With time, tears become more complex, degenerative and less amenable to repair(79–84).
- 5.7.2 Repairing the meniscus at the same time as ACL reconstruction improves the success rate of the repair(85).
- 5.7.3 Significant lateral meniscal injury, particularly bucket handle and meniscal root tears are associated with a high grade pivot shift(86,87).
- 5.7.4 Medial meniscus posterior horn injury can predispose to premature ACL graft failure(88).
- 5.7.5 Repair of repairable meniscal tears (compared to meniscectomy at the time of ACL reconstruction) may result in improved subjective knee pain and function(89), objective knee scores and reduce the progression of radiographic features of osteoarthritis(90–92).
- 5.7.6 Small, stable partial thickness fissures in the posterior horn of the lateral meniscus can usually be left untreated(93).
- 5.7.7 Patients presenting with ACL injury and a locked knee secondary to a bucket handle meniscus tear may develop fixed flexion deformity if untreated. The meniscus tear is more likely to be repairable if reduced early; the ACL may be reconstructed concomitantly or as a delayed procedure following the meniscus repair, when knee swelling has reduced and range of movement has been recovered. In a staged approach, the knee should be protected in a ROM brace. Concomitant repair of meniscal tears with ACL reconstruction improves meniscal healing (85).

## **5.8 Articular cartilage injury.**

5.8.1 Articular cartilage injuries range from minor fissuring to full thickness chondral or osteochondral loss and fracture(38).More damage is seen in chronic cases(83).

5.8.2 More urgent intervention may be required to facilitate fixation of large osteochondral injuries or removal of loose bodies that are hampering pre-habilitation.

5.8.3 Treatment options for chondral defects depend on: chronicity, patient age, defect size and location. Surgeons undertaking ACL reconstruction should be familiar with these options. Treatment should be based on published outcome data for various techniques as well as guidance from the UK Cartilage Consensus statement(94).

## **6.THE JUVENILE ACL**

### **6.1 Introduction**

6.1.1 The incidence of juvenile ACL rupture is rising with juvenile patients now representing a small but significant portion of the work of many knee surgeons(95–97). One recent study reports 29 fold increase in the rates of paediatric and adolescent ACL reconstruction in UK over the last 20 years(98).

6.1.2 Injury prevention programs can reduce the incidence of ACL rupture in some groups and their use in schools should be encouraged(99–101).

6.1.3 Clinical and radiographic assessment can be difficult in juvenile patients with injuries in continuity disproportionately common.

6.1.4 Treatment presents a therapeutic dilemma. The consequences of the injury include progressive intra-articular pathology & instability and unfortunately there are risks associated with all types of treatment.

6.1.5 The current consensus is that in symptomatic patients the risks associated with conservative treatment exceed those associated with surgery(102).

6.1.6 Results following conservative or operative management in this group are less favourable compared to the adult population.

6.1.7 Due to the challenges posed by caring for this specific group of patients, BSCOS and BASK have initiated a national steering group to provide more specific evidence-based guidance on care of these patients. While this process is ongoing, the following recommendations serve as interim guidance, which will be carried forwards and expanded upon within the full joint BSCOS/BASK joint committee practice framework.

## **6.2 Conservative Treatment**

6.2.1 Conservative treatment includes bracing, activity modification and structured rehabilitation and is very difficult to provide effectively. No studies directly compare conservative vs surgical treatment but meta-analysis reveals multiple trends favouring early surgery with patients treated surgically experiencing less instability and more able to return to previous levels of activity(103).

## **6.3 Surgical Treatment**

6.3.1 Some injuries, including repairable meniscal tears, require early surgical treatment and in these cases, reconstruction should also be undertaken.

6.3.2 Symptomatic instability is the key indication for reconstruction and delay in treating symptomatic patients increases the risk of developing further meniscal & chondral pathology(104–106).

6.3.3 Surgical reconstruction is broadly of two types, either Transphyseal or Physeal Sparing. Although repair has been dismissed in the past some surgeons have also recently reported using augmented repair for proximal avulsions in young patients(107).

6.3.4 Transphyseal reconstructions have been reported on many hundreds of juvenile patients in all Tanner stages and with the largest case series reporting good or excellent results using hamstring tendon autograft. This technique is currently the most widely used in juvenile patients(108–114).

6.3.5 Physeal Sparing Extraphyseal reconstructions are variants of the MacIntosh & Micheli procedures. Relatively small numbers of these non-anatomic procedures have been reported(115–117).

6.3.6 Physeal Sparing All Epiphyseal reconstructions become increasingly difficult to perform in younger children with smaller physes. Surgeons regularly performing all inside reconstruction may be better equipped to use this technique(118–120).

6.3.7 Hybrid techniques combining Transphyseal tibial and Physeal Sparing All Epiphyseal femoral tunnels have also been reported(121).

6.3.8 The lack of consensus on surgical treatment has prompted the Paediatric Anterior Cruciate monitoring initiative (PAMI) recently announced by ESSKA(122,123).

## **6.4 Results of Surgical Treatment**

6.4.1 Functional results following juvenile ACL reconstruction are generally good and in some large transphyseal series excellent. Juveniles report better functional outcomes than adults and are more likely to return to sport(111,114).

## **6.5 Iatrogenic Physeal injury**

6.5.1 Direct physeal injury is caused by the drilling & reaming of bone tunnels and the associated heat generated. Larger tunnels are more damaging as are tunnels that are very oblique or very peripheral. Direct injury causes bone bridging with the presence of a soft tissue graft thought to be protective. Indirect injury includes physeal compression due to over tensioning, and vascular phenomena which can cause either growth arrest or overgrowth(124–126).

6.5.2 The greater the potential for growth the greater the consequences of growth arrest and the treatment of prepubescent patients in Tanner stage 1 & 2 gives the most cause for concern. Assessing the patient with their parents in clinic provides useful information and surgeons should consider making formal assessments of bone age and maturity when undertaking surgical treatment, including radiologic calculation of skeletal age.

## **6.6 Growth Disturbance**

6.6.1 Growth disturbance does occur following reconstruction and historically its incidence may have been under reported(123,127).

6.6.2 Growth disturbance has been reported following Transphyseal and all of the available Physeal Sparing reconstruction techniques. Deformities include leg length inequality and angulation usually into valgus or recurvatum. Evolving deformity may not be apparent to the patient and may not be obvious clinically(128–131).

6.6.3 The incidence of growth disturbance is unknown but it is currently believed to be uncommon. The two largest Transphyseal series report no significant growth disturbance. Few studies accurately assess for growth disturbance. A meta-analysis found growth disturbance in only 19 of over 900 cases at risk with higher rates following Physeal Sparing than Transphyseal reconstruction. A case series using long standing radiographs reported a

4% incidence of coronal deformity exceeding 2 degrees (but none exceeding 5 degrees) at 2 yrs(108,109,111,112,114,132).

6.6.4 Surgeons performing ACL reconstruction in juveniles should undertake regular clinical review and should consider arranging regular radiographic review, including pre and postoperative long standing radiographs, until their patients have reached maturity(102,131).

6.6.5 The involvement of Paediatric Orthopaedic Surgical colleagues facilitates the monitoring for and treatment of growth disturbance.

## **6.7 Graft rupture and Contralateral ACL Rupture**

6.7.1 Rates of graft rupture and contralateral ACL rupture are disproportionately high in juvenile patients with young age, return to sport and family history, the most important factors associated with an increased risk of re-injury. Around 1/5 juvenile patients will sustain an ACL graft rupture, and around 1/5 a contralateral ACL rupture within 5yrs of surgery. Altogether around 1/3 will eventually sustain a re-injury with most of the graft ruptures occurring within 2 years and most of the contralateral ACL ruptures between 3 & 5 years(112,133–138).

6.7.2 Although in Europe around 90% of juvenile ACL reconstructions are performed using hamstring tendon autograft some evidence suggesting that re-rupture rates in younger patients may be lower using patella tendon is emerging. Re-rupture rates using irradiated allograft in juveniles are high but one small case series using fresh allograft donated by parents has reported good results and low rates of graft rupture at 2 years(95,133,139–143).

## **6.8 Rehabilitation**

6.8.1 Graft maturation continues for at least 12 months with neuromuscular performance impaired for up to 18 months and a conservative rehabilitation regime should generally be employed in juvenile patients. Although some authors recommend postoperative bracing it was not used in the largest Transphyseal series and there is no evidence supporting its routine use(144).

## **6.9 Conclusion**

- 6.9.1 In symptomatic patients the benefits of surgical treatment outweigh the associated risk.
- 6.9.2 Assessments of bone age and maturity should be made when undertaking surgical treatment.
- 6.9.3 When performing ACL reconstruction in juveniles, surgeons should use the technique with which they are most familiar and perform the best reconstruction that they can whilst respecting the physis.
- 6.9.4 Surgical technique should be modified with longer less oblique tunnels of 7-8mm or less in diameter, very slow reaming avoiding heat, and care taken to avoid injury to the tibial tubercle apophysis and the femoral perichondrial ring. Bony debris should be cleared from tunnels, and if used bone blocks and screw threads should not be placed across the physes.
- 6.9.5 Functional outcomes are usually good following reconstruction.
- 6.9.6 The risk of significant growth disturbance remains relatively low.
- 6.9.7 The use of pre & postoperative long leg standing radiographs facilitates the detection of evolving deformity.
- 6.9.8 The involvement of Paediatric Orthopaedic Surgical colleagues facilitates the monitoring for and treatment of growth disturbance.
- 6.9.9 Re-injury rates are disproportionately high in juveniles and so patients and their families should be counselled accordingly.
- 6.9.10 For further guidance surgeons should refer to the BSCOS/BASK Steering committee report on the Child and Adolescent Knee.

## **7-PRE-OPERATIVE ASSESSMENT AND CONSENT**

- 7.1 A systematic multidisciplinary pre-operative assessment is recommended as good practice. This assessment should include the presence of allergies, including latex, and co-morbidities e.g. diabetes mellitus, both for patient management and list planning. The risk for deep venous thrombosis should also be determined and prophylactic treatment based upon local and national treatment guidelines.
- 7.2 Patients should be swabbed for Methicillin Resistant Staph. Aureus and treated prior to the day of surgery (145).

- 7.3 The social support available for the patient should be ascertained to determine whether the procedure should be performed on a day case or in-patient basis.
- 7.4 The knee should be assessed to ensure full extension has returned, the knee is ideally free from an effusion and quadriceps activation and rehabilitation has been commenced prior to reconstruction(146–148)
- 7.5 Ideally the consent process should have started in the clinic, reaffirmed at the pre-operative assessment and continue on the day of surgery. Guidance on consent process has been given by the Royal College of Surgeons(149).
- 7.6 Operative and non-operative options should be discussed. The likely outcome of each including the benefits and limitations should be discussed and recorded(10,32,150). The general risks of surgical intervention together with the specific risks of reconstructive surgery should also be discussed and this conversation documented(151–157). Risks of persistent instability and reinjury, including to the opposite leg should be discussed(133,158,159). Patients should be advised on timing of return to sports, which should be specific to the individual (160–162), but should rarely be less than 9 months before surgery due to higher risk of reinjury (160).
- 7.7 Graft selection should be confirmed whether autograft, allograft or synthetic ligaments are to be used. This should include the benefits, complications and risks of the preferred graft. The frequency of complications together with their management and implications should be included in part of the consent process(151–153).
- 7.8 The management of chondral and meniscal pathology should also be discussed together with the implications on post-operative recovery. This allows the patient the opportunity to reflect on these discussions and ask further questions.
- 7.9 The National Ligament Registry ([www.uknlr.co.uk](http://www.uknlr.co.uk)) should be discussed with all patients including the benefits and importance of participation. Patients should be registered and consent obtained for inclusion of their data. Hospitals should facilitate the accurate recording of surgical procedures and patients' outcome by providing appropriate clerical and IT support.



7.10 Imaging should be reviewed as during the pre-assessment and if radiographs and scanned images were brought with the patient, they should be ensured that images are uploaded to the imaging system at the operative site. These can then be available to view in operating theatre during surgery.

## **8 THE ADMISSION TO HOSPITAL**

8.1 The patient should be fully aware of the intended procedure prior to admission. The benefits, risks and potential complications of ACL reconstruction should be repeated. Emphasis should be given to the role of the pre and post-operative rehabilitation programme.

8.2 The surgical site should be marked in an area which is still visible after draping.

8.3 The patient must reconfirm consent to the operating surgeon or a suitably qualified deputy

8.4 The procedure can be performed on a Day Case basis, with appropriate domiciliary support.

## **9 HOSPITAL FACILITIES REQUIRED FOR THE OPERATION**

9.1 Primary ACL reconstruction operations are best carried out in hospitals where knee ligament surgery is regularly performed and adequate numbers of trained nurses and the skills of Professions Allied to Medicine are available.

9.2 The potential for cross-infection should be reduced to a minimum. Patients should be admitted to and nursed on elective orthopaedic or day case wards that are staffed by a team experienced in the care of patients who have undergone ligament reconstructions.

9.3 The use of ultra-clean air theatres is still considered to be best practice for units performing any surgery with the implantation of foreign material or open joint surgery, though controversy exists on efficacy of laminar flow theatres(163–165) There is, however, no published evidence specifically in respect of ACL reconstruction to support this.

9.4 The operating theatre should be dedicated to clean elective orthopaedic surgery or joint

reconstruction. Shared facilities with other clean surgical disciplines is an acceptable practice, when using ultra clean air, but data supporting this practice are not available.

9.5 The surgeon must have trained assistance during the operation, and a trained scrub nurse fully familiar with the required complex instrumentation is mandatory. In the absence of junior staff, additional Nursing assistants or specifically trained Surgeon's assistants must be available.

9.6 Endoscopic ACL reconstruction is the gold standard. A fully functional arthroscopic stack camera system and fluid management system must be available. Power shaving system or radio-frequency ablation systems must also be available according to surgeons' preferences.

9.7 A full range of specialist implants and instruments suitable for both hamstrings and patellar tendon reconstructions must be available before the start of each case.

9.8 Specialist instruments and devices should be readily available to manage concomitant meniscal or chondral pathologies.

9.9 Image intensifier may be necessary for some cases such as paediatric physeal sparing ACL reconstructions.

9.10 Dynamic or static knee braces may be required in selected cases.

9.11 In units with several surgeons performing ACL reconstructions, it is desirable to form a consensus for a single instrument system and implants to be used. This is to enhance inventory and reduce cost. It will also help with theatre staff skill level and training.

9.12 Appropriate impenetrable clothing and drapes are essential.

## **10 ANAESTHESIA**

10.1 ACL reconstruction will usually be carried out as an elective day case procedure under general anaesthesia, but spinal anaesthesia can be an alternative if deemed appropriate. Royal College of Anaesthetists have provided guidelines on standards of care for the provision of daycase

surgery (166). Recently, the Society for Ambulatory Anesthesia, in America have issued evidence based recommendations on perioperative pain management for ACL reconstruction(167).

10.2 Pre-assessment for anaesthesia- Specific arrangements and processes for pre-anaesthesia assessment will differ in detail between hospitals but the number of hospital visits for the patient should be minimised. Where possible, anaesthesia assessment should be synchronised with the orthopaedic clinic visit.

10.2.1 The majority of patients will be fit young adults requiring no specific pre-operative tests.

10.2.2 Pre-assessment aims to identify and delineate any medical co-morbidity (e.g. asthma, diabetes, hypertension, epilepsy) that may require optimisation, stabilisation or preparation prior to surgery.

10.2.3 Problems that may influence the selection of anaesthetic technique and choice of post-operative analgesia as well as suitability for day surgery are also best identified at the pre-assessment visit.

10.2.4 There may be a need to prioritise specific patients (such as diabetics) first on the operating list and planned overnight hospital admission may be necessary.

10.3 Anaesthesia- Consent for anaesthesia is usually obtained verbally and explanations of the risks of anaesthesia will be documented by the anaesthetist along with ASA (American Society of Anesthesiologists) grade.

10.3.1 Thrombo-embolic risk and requirement for prophylaxis will be evaluated for each patient in collaboration with surgical colleagues.

10.3.2 Fasting times prior to anaesthesia should follow local guidelines with clear fluids usually being permitted until 2 hours pre-op as a maximum(168). 250 mls of water should be encouraged prior to the start of the theatre list to avoid dehydration.

10.3.3 Local safety checklists and WHO guidelines should be followed

10.3.4 General anaesthesia for ACL reconstruction does not routinely require muscle relaxant or intubation. Inhalational or intravenous anaesthetic agents along with analgesics (opiates, NSAIDs & other adjuncts) and anti-emetics are deployed according to the anaesthetist's preference and patient's medical status.

10.3.5 Antibiotic prophylaxis will need to be administered prior to inflation of the tourniquet.

10.3.6 Multimodal analgesia inclusive of 2 or more non-opioid analgesics (e.g. Paracetamol

and NSAIDs or Cox-2 inhibitors) should be used for perioperative pain relief and to limit postoperative opioid use(167,169). Intraoperative opioid sparing drugs (e.g. Clonidine and MgSO<sub>4</sub>) can aid in postoperative pain control and same day discharge.

10.3.7 Administering a preoperative single 600-mg dose of gabapentin may decrease both pain intensity and opioid consumption in patients undergoing arthroscopic ACL reconstruction(170).

10.3.8 A nerve block can be used to provide post-operative pain relief especially in patients unable to tolerate systemic analgesics(171). However, with local infiltration analgesia and multimodal analgesia, most patients can be managed without(167). Recommendations for patient explanation, consent and safety checks should be followed

10.3.9 Femoral Nerve block (FNB) provides good quality analgesia but concurrent motor weakness can delay full weight bearing and there are concerns about persistent quadriceps deficit (167,172,173). There is lack of evidence of benefit to femoral nerve block when used in the setting of local instillation analgesia(174).

10.3.10 Saphenous nerve block in adductor canal has been shown to be equally or more effective than femoral nerve block, without the risk of quadriceps weakness(175–179). Again, there is limited evidence of its benefit over local anaesthetic infiltration (180–182).

10.3.11 Local anaesthetic infiltration following ACL reconstruction has been shown to be equally effective in controlling postoperative pain and reducing opioid use, without any risks (167,174,180,182,183). However, for patients with risk factors such as tolerance to opioids or preoperative chronic pain, either adductor canal block or FNB for ACL reconstruction should still be strongly considered and is consistent with properly practiced evidence-based medicine(184).

10.3.12 There are some concerns about the intraarticular toxicity of local anaesthetics (185,186)which can be addressed by periarticular and pericapsular infiltration(187). When hamstring graft is used, harvest site should be infiltrated with local anaesthetic(188).

10.3.13 There is limited evidence to support the use of indwelling catheters(187)

#### 10.4 Post-op recovery & discharge.

10.4.1 Standard practice for day case anaesthesia, recovery and analgesia including nurse led discharge should be followed(166). This includes opiate sparing analgesia to take home

(paracetamol and NSAID) to minimise post-operative nausea & vomiting (PONV).

10.4.2 Warnings about impaired co-ordination and balance for at least 24 hours post-op should be given (particularly regarding stairs and with crutches) and when stronger analgesics are prescribed.

10.4.3 Next day telephone follow-up by day unit staff can provide useful audit data and patient reassurance.

## **11 THE SURGEON**

11.1 All patients should be admitted under the care of a consultant orthopaedic surgeon who has a specialist interest in the surgical management of the anterior cruciate deficient knee.

11.2 The surgeon may delegate all or part of the procedure to a trainee or another surgeon who is developing their skills in ACL reconstruction.

11.3 The supervising consultant may or may not be scrubbed at the operating table when teaching a surgeon in training. That decision should be made jointly between the trainer and trainee and would depend on the experience of the latter.

11.4 The lead surgeon undertaking ACL reconstruction should have received appropriate and adequate training and have sufficient experience to deal with the case including the development of unexpected intra-operative complications.

11.5 The issue of “minimum numbers” is a debatable and controversial topic. There is increasing evidence in other areas of surgery that better outcomes are achieved by high volume surgeons. Vice versa, poorer outcomes are associated with low volume surgeons. No absolute minimum numbers are recommended but it would be considered desirable for a surgeon to be performing ten or more ACL reconstructions per year.

11.6 The theoretical and practical skills of a surgeon performing primary ACL reconstructions must be maintained by relevant Continuing Professional Development (CPD). Such evidence should be presented annually at the surgeon’s appraisal.

11.7 It is expected that the surgeon will be registered with the National Ligament Registry (NLR)

and that every operation is centrally databased. Any clinician feedback data generated from the NLR must be included and discussed at the surgeon's annual appraisal.

11.8 The surgeon should preferably be a member of one of the relevant specialist societies (BASK British Association for Surgery of the Knee) or BOSTAA (British Orthopaedic Sports Trauma and Arthroscopy Association).

## **12 PROPHYLAXIS AGAINST VENOUS THROMBOEMBOLISM (VTE)**

12.1 The risk of deep venous thrombosis following ACL surgery remains very low(155–157,189,190), therefore pharmacological VTE prophylaxis is not indicated as a routine. However, all patients should be risk assessed upon admission to hospital. In high-risk cases, and when the reconstruction is combined with posterior cruciate reconstruction, thromboprophylaxis should be considered.

12.2 Patients who had ACL reconstruction considered as being at increased risk of VTE when:

12.2.1 The surgical procedure takes more than 90 minutes

12.2.2 The patient or first-degree relative with a history of VTE

12.2.3 The patient uses hormone replacement therapy or estrogen-containing contraceptive therapy

12.2.4 They have varicose veins with phlebitis

12.2.5 The patient is obese (body mass index [BMI] over 30 kg/m<sup>2</sup>)

12.2.6 They have travelled recently on long haul flight.

12.2.7 They aged  $\geq 35$  years(155,189)

12.3 Patients can reduce the risk of VTE by starting the rehabilitation program early, staying hydrated and becoming more mobile. Mechanical prophylaxis like calf pumps should be used intraoperatively and in recovery following surgery, before patient becomes mobile.

## 13 SURGICAL PROCEDURE

13.1 An examination under anaesthetic must be performed to take into account the degree of anteroposterior and rotational laxity as well as any other associated injuries. The findings may influence the need to proceed to adjunctive procedures.

13.2 Tourniquets are frequently used but are neither mandatory nor contraindicated. The tourniquet may be inflated for part of the procedure, or for all of it.

13.3 The skin should be prepared with an appropriate alcohol based solution(191).

13.4 The surgical procedure of ACL reconstruction can take a variety of forms. Whether ACL reconstruction is undertaken arthroscopically (most common) or open (increasingly rare), arthroscopic examination of the joint to deal with any other associated intra-articular injuries is necessary. Most surgeons undertake the entire procedure arthroscopically, although incisions are needed for graft harvest, for femoral tunnel drilling or fixation in some techniques, and for extra-articular tenodesis when that is undertaken simultaneously.

13.5 The ACL is typically reconstructed using autogenous tissue but allograft or synthetic tissue may be used. The type of graft to be used should be discussed with the patient pre-operatively. The choice should be tailored to the individual and data entered into the National Ligament Registry. Surgeons should be familiar and proficient in the technique chosen.

13.6 The most common autografts being used include bone-patellar tendon-bone (BPTB), hamstring tendon (HT), and quadriceps tendon (QT). They all have been shown to have good outcome for majority of patients(151,192,193). Hamstring tendon might have a slightly higher re-tear rate when compared with BPTB (139,140,151) with main complication being of damage to the infrapatellar branches of the saphenous nerve(194). However, BPTB has a higher rate of anterior knee and kneeling pain(194) with suggestion of greater risk of osteoarthritis (195,196). Quadriceps graft have been shown to have less harvest site morbidity than BPTB graft with good functional outcome(193,197,198) but some studies suggest higher failure rates(199). Long term data from large studies on quadriceps tendon graft is awaited.

13.7 Allografts include hamstring grafts, Achilles grafts, patellar tendon grafts and other grafts

sourced from a number of other lower limb tendons. Surgeons should understand the source of the graft and the technique used for cleansing and storage(200,201). Patients should be aware of potential risks of infection and premature failure with some graft materials. Allograft is a viable option for revisions and primaries in patients greater than 35 years old as they avoid donor site morbidity; however, re-tear rate increases significantly in younger patients(202–204).

13.8 Graft choice should ultimately be decided upon based on surgeon comfort, experience and individual patient characteristics and should be one part of a larger conversation with each individual patient.(151)

13.9 Synthetic ligaments are not currently recommended for routine primary intra-articular reconstruction.

13.10 Wrapping of graft in Vancomycin soaked swab (5mg/ml), prior to implantation, has been shown to significantly reduce infection rates in ACL reconstruction surgery to approaching 0%(205–211). Experimental studies suggest the use of 5mg/ml Vancomycin soaked swab for 20 minutes decontaminates graft without affecting its biomechanical properties(212–214). No deleterious effects on surgery outcome or increased graft failure rates, have been reported in currently available literature(205,206). However, Vancomycin has not been compared against other antibiotics and wider issues about future drug resistance need to be considered(206,215)

13.11 Partial ACL injuries may also be augmented with allograft, autograft or synthetic tissue and there are procedures to 'repair' the ACL in selected cases for acute injuries(162,216–221). Repair techniques are currently experimental and should be undertaken as part of studies and within clear governance processes in the organisations where those procedures take place(222).

13.12 An ACL reconstruction requires tunnels in the femur and in the tibia. These tunnels can be drilled outside-in or inside-out. There are many theories regarding optimal tunnel placement(223–228). The anatomy of the knee needs to be clearly visualised in order to allow the surgeon access to their planned tunnel entry points. The use of shaver and electro-cautery systems is helpful.



- 13.13 The femoral tunnel is typically drilled through a transportal technique, although it may be drilled through a transtibial technique. The latter technique may constrain the position of the femoral tunnel(229,230).
- 13.14 The femoral tunnel is usually placed in an anatomical or near anatomical position. This is an area of great debate but is generally between 9:00 and 11:00 or 1:00 and 3:00(223–227). If the femoral tunnel is too posterior, it risks blowing out the posterior wall. If the femoral tunnel is too anterior, deep flexion is restricted and graft failure may ensue when motion is regained.
- 13.15 The tibial tunnel is typically placed in the middle third of the tibia within the ACL footprint. If the tunnel is too anterior, it leads to impingement and loss of extension. If the tunnel is too posterior, it impinges on the PCL(228).
- 13.16 Preservation of native tissue is encouraged wherever possible, although the data to support this is limited. Extra care should be taken when preserving the tibial stump as this may obscure tibial tunnel position and increase the risk of impingement(218).
- 13.17 During the procedure the surgeon should test that the graft position achieved does not result in loss of motion or in impingement.
- 13.18 Fixation can be undertaken in several ways. Suspensory fixation has gained popularity, but screw and cross-pin fixation is also possible on the femoral side. Fixation on the tibial side may be within the bone but can also be suspensory on the surface of the tibia. Once the graft has been fixed on one side, the knee should be cycled to ensure there is no impingement or excess graft motion, and it should then be fixed on the other side. Surgeons should be aware of the intricacies of fixation techniques that are used(231,232). Fixation technique used should allow for immediate knee movement
- 13.19 Both the graft harvest site and arthroscopic portals should be closed in an appropriate manner and compressive dressings applied.
- 13.20 **Lateral tenodesis.** Clinical evidence is currently emerging to support use of lateral extra-articular procedures, in certain indications, as an augmentation to the ACL reconstruction. Appropriate indications may include, patients younger than 25 years; patients with generalized

ligamentous laxity or genu recurvatum; patients with high grade pivot shift or in revision ACL reconstructions. Such reconstructions may have higher early morbidity but can reduce long term graft failure rates(233–239).

## **14 POSTOPERATIVE CARE AND FOLLOW UP**

14.1 Analgesia-A locally agreed regime should be adopted in liaison with anaesthetic colleagues. This should include regular analgesia as well as adequate provision for breakthrough pain.

14.2 An Xray in the post-operative period is useful for confirming the position of the fixation devices and bone tunnels. Post-operative imaging is recommended for all patients where there is any clinical concern about symptoms or progress with rehabilitation.

14.3 **Bracing**-There is little evidence to support the routine use of brace following ACL reconstruction(240–248). Some data suggest that functional bracing may protect reconstructed graft during stressful activities(249). Occasionally surgeons may use a brace, at their discretion. Factors that may influence a consultant’s decision to brace include:

14.3.1 Combined procedure such as Meniscal repair or Micro fracture.

14.3.2 Use of Femoral nerve block.

14.3.3 To facilitate day case discharge.

14.3.4 Preference towards slow early mobilisation.

14.3.5 In adolescents, where use of brace has been shown to reduce the risk of graft re-tear(250).

14.3.6 To protect the graft on return to sport(246).

14.4 Weight bearing-All patients should be encouraged to fully weight bear as comfort allows, unless weight bearing is contraindicated by a concomitant procedure such as Micro fracture, high tibial osteotomy etc.

14.5 Wound care-Clear instructions should be given to the patient and all allied staff, as to the operating surgeons wishes in relation to timing of dressing changes, wound inspections, and removal of any stiches.

14.6 Follow up-The follow-up arrangements vary widely across the UK. At a minimum, patients

should be followed up and reviewed within the orthopaedic service at the following post-operative time points:

14.6.1 - Approximately 2 weeks. Wound review, surveillance for complications, facilitation/referral for rehab.

14.6.2 - Approximately 6-12 weeks. Assessment of early rehab goals, resolution of effusion, regaining range, early strength.

14.6.3 - Approximately 24-30 weeks. Assessment of late rehab goals, transition to sport.

14.6.4 Follow up may be conducted by the clinical team or trained allied healthcare professionals as part of a wider multi-disciplinary team. In the later stages the patient's 'Follow up' may be conducted within the rehab setting and under the care of experienced and specialist physiotherapists.

14.7 Commencement of Physio-Patients should be commenced upon an ACL specific rehabilitation programme following their post-operative follow up appointment. Please refer to the rehabilitation section of these guidelines.

## **15 NATIONAL LIGAMENT REGISTRY**

15.1 Injury to the anterior cruciate ligament (ACL) is common but the incidence of ACL injury in the UK is unknown, the number of surgeons undertaking such surgery (and what volumes) is not clear and patient outcomes are only reported in single surgeon or small unit studies. If we apply the incidence of 1:80,000 (Swedish ACL registry report 2018)(251) to the UK population it would suggest that approximately 54,000 ACL injuries occur per annum in the UK. If 50% of these patients require surgery this would result in 27,000 primary ACL reconstructions each year, 3.5 times greater than the number of unicompartmental knee replacements recorded on the National Joint Registry(252)

15.2 The increased risk of degenerative joint disease following ACL rupture is well known but there is no clear understanding of the rate of repeat / revision surgery following ACL reconstruction, which could be as high as 20%. Arthroplasty publications suggest poorer outcomes with lower volume surgery, but it has yet to be determined whether similar trends occur with soft tissue knee surgery.

15.3 Our aim is to perform interventions which are the best and safest for our patients and the

recording of patient outcomes should be an essential component of patient care.

15.4 The National Ligament Registry (NLR) was established in 2013 to initially capture data on primary ACL reconstruction. It comprises an online data collection system which automatically contacts patients by email at certain time intervals and provides a secure link for them to securely enter the various PROMs. The selected outcome measures are the Knee injury and Osteoarthritis Outcome Score (KOOS), subjective International Knee Documentation Committee (IKDC), Euroqol (EQ5D) and the Tegner activity score. These scores allow comparison and communication with existing ACL registries as well as allowing potential 'generic health benefit' comparisons to other non-Orthopaedic procedures. The registry should provide patients with an element of 'quality assurance' which extends beyond patient functional outcome to surgical numbers, patient selection, rehabilitation, prostheses and surgical techniques.

15.5 The main roles of the NLR are to:

15.5.1 Encourage the development of better surgical practice for our patients.

15.5.2 Provide a framework for surgeons to collate and audit their results, satisfying the modern requirements of appraisal and revalidation.

15.5.3 Record all aspects of each specific injury and intervention, including primary ligament repair, differing graft options and fixation types, allowing equitable comparison.

15.6 The website ([www.uknlr.co.uk](http://www.uknlr.co.uk)) has been developed for both patients and professionals and in time we hope will be the 'go to place' following ACL injury. It contains patient and surgeon information videos and a mobile platform has been developed for use with tablets and smart phones. There is also a 'map' of NLR registered surgeons which will enable patients to see where surgeons who record their ACL reconstruction data on the NLR are based.

15.7 The NLR is independent of both Government and Private Medical Insurance company support and receives its' financial support from industry partners involved in various aspects of ACL reconstruction surgery.

15.8 Big collections of data are powerful provided that such data is interpreted, presented and incorporated in a responsible, professional way. The key to the success of the registry is the involvement of all stakeholders and it is important that patients, surgeons and industry are

involved, feel valued and benefit from the process.

15.9 Annual reports are published to coincide with the BASK annual meeting and are available to view or download as a PDF from the website.

15.10 Compliance remains a challenge to all registries and at the time of writing the NLR is not yet a 'mandated' Orthopaedic registry. However, the submission of data to all established registries is supported by the BOA and the NLR is also strongly supported by BASK. Discussions are ongoing about NLR involvement in the BOA registry framework, known as TORUS.

15.11 With the increasing requirements for surgical outcome data the NLR is a ready to use online data collection system which has the functionality to provide information at surgeon level (appraisal/revalidation), unit level and national level. It is overseen and coordinated by surgeons and will ultimately provide the information we require to continue treating patients with this type of intervention in the best way we can.

15.12 It is recommended that trusts would provide administrative support to help input patient data into National Ligament Register and all surgical reconstructions of the ACL would be documented in the ligament registry.

## **16 CONSERVATIVE MANAGEMENT, REHABILITATION AND RETURN TO SPORT OF ACL INJURY (RECONSTRUCTED) PATIENT**

### **16.1 Pre-operative management**

16.1.1 Knee function at time of surgery has been shown to be a significant factor in predicting outcome(253). Preoperative quadriceps weakness (greater than 20% difference between sides) has been shown to predict poor quadriceps strength & low self-reported function after surgery(146,254). It would be recommended that the patient engage in a lower limb, especially quadriceps strengthening programme prior to surgery.

16.1.2 A restriction of range of movement pre-operatively has been shown to significantly limit the progression of post-operative rehabilitation (146,147), if the patient does not have full range of movement pre-operative it may prove useful to undertake a course of physiotherapy to regain range of movement.

## **16.2 Management of the immediate post-operative phase**

- 16.2.1 Persistent quadriceps lag on straight leg raise has been shown to indicate an inability to actively fully extend the knee. If this is not achieved by week 5 post-operation this would be considered a predisposing factor for significant quadriceps weakness at 6 months post operation. This activation failure of the quadriceps is likely to serve as a major barrier to rehabilitation limiting rehabilitation progress(255). Early activation of quadriceps can be achieved through exercise and/or neuromuscular electrical stimulation. Early joint motion is beneficial in order to avoid capsular contractions, reduce swelling and pain. Even small losses of knee extension (3-5 degrees) appear to adversely affect subjective and objective outcome markers later in the rehabilitation phase(248). Early quadriceps activation and full range motion should be encouraged and where necessary support with an appropriate physiotherapy programme.
- 16.2.2 Abnormal gait patterns have been associated with low patient satisfaction and decreased functional performance(256).These gait abnormalities also often become further exaggerated when the patient returns to running (257). It is important then to retrain the patients gait in the immediate post-operative period.

## **16.3 Rehabilitation phase**

- 16.3.1 The retraining of postural and movement control requires the incorporation of both static balance exercises along with dynamic balance and movement tasks (247). The regaining symmetry in relation to limb alignment control during limb loading activities after ACLR may significantly reduce their potential for future ACL injury (258). Progressive strengthening programmes for key muscles (quadriceps, hamstrings & gluteal muscles) is required to prepare the patient for controlled load acceptance and subsequent return to sport (247). It is not uncommon to have significant deficits in rate of force development in patients post ACLR which may impact on the functional performance and affect the ability to perform sport specific speed and agility based tasks(259,260). There is a strong relationship between cross over hop performance and functional outcome(261) correlating significantly to IKDC subjective and KOOS questionnaire scores(262). Once the patient has sufficient strength and static-dynamic balance, they need to have rehabilitation exercises targeting both rate of force development and plyometric ability.

## **16.4 ACLR and return to sport (RTS)**

- 16.4.1 There are now guidance available guiding assessment criteria to be used before allowing patient to return to unrestricted sports activities (161,263–266). RTS is a continuum comprising three elements: return to participation, return to sport and return to performance(266). Decision to return to sport should be criteria based, taking into consideration- physical factors relating to the knee; psychological factors including fear of reinjury and social factors; while being tailored to the specific sport. To assess readiness to return to play and the risk for reinjury, a battery of tests, including strength tests, hop tests and measurement of movement quality, should be used(161). These tests should include assessments of direction changes and reactive agility tests and the assessment of psychological readiness to RTS using an instrument such as the ACL-Return to Sport after Injury scale(267)
- 16.4.2 Gradual and sequential introduction of sport-specific training can be used as functional tests that include an element of protected reactive decision-making.(266)
- 16.4.3 Return to sport 9 months or later after surgery and with symmetrical quadriceps strength prior to return substantially reduces the reinjury rate(160,161)

#### **16.5 Conservative management of ACL injured patient:**

- 16.5.1 Here the goals would be to minimise any functional instability of the knee and mitigate against future degenerative changes within the knee. These two goals are not mutually exclusive. Improving strength, dynamic and static balance will improve knee stability and reduce the tendency towards co-contraction and increased shear and compressive forces which add increased stress onto the articular surfaces. The global management strategy is very similar to that for ACL reconstruction.

## BIBLIOGRAPHY

1. The National Ligament Registry [Internet]. [cited 2020 Jan 13]. Available from: [www.uknir.co.uk](http://www.uknir.co.uk)
2. Ardern CL, Taylor NF, Feller JA, Webster KE. Fifty-five per cent return to competitive sport following anterior cruciate ligament reconstruction surgery: an updated systematic review and meta-analysis including aspects of physical functioning and contextual factors. *Br J Sports Med* [Internet]. 2014 Nov [cited 2019 Sep 21];48(21):1543–52. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25157180>
3. Parwaiz H, Teo AQA, Servant C. Anterior cruciate ligament injury: A persistently difficult diagnosis. *Knee* [Internet]. 2016 Jan [cited 2019 Sep 21];23(1):116–20. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26552783>
4. Lau BH, Lafave MR, Mohtadi NG, Butterwick DJ. Utilization and cost of a new model of care for managing acute knee injuries: the Calgary acute knee injury clinic. *BMC Health Serv Res* [Internet]. 2012 Dec 5 [cited 2019 Sep 21];12(1):445. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23216946>
5. Perera NS, Joel J, Bunola JA. Anterior cruciate ligament rupture: Delay to diagnosis. *Injury* [Internet]. 2013 Dec [cited 2019 Sep 21];44(12):1862–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24012476>
6. Smith SD, LaPrade RF, Jansson KS, Årøen A, Wijdicks CA. Functional bracing of ACL injuries: current state and future directions. *Knee Surgery, Sport Traumatol Arthrosc* [Internet]. 2014 May 27 [cited 2019 Sep 21];22(5):1131–41. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23624655>
7. Filbay SR, Grindem H. Evidence-based recommendations for the management of anterior cruciate ligament (ACL) rupture. *Best Pract Res Clin Rheumatol* [Internet]. 2019 Feb [cited 2019 Sep 21];33(1):33–47. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/31431274>
8. Beaufils P, Hulet C, Dhénain M, Nizard R, Nourissat G, Pujol N. Clinical practice guidelines for the management of meniscal lesions and isolated lesions of the anterior cruciate ligament of the knee in adults. *Orthop Traumatol Surg Res* [Internet]. 2009 Oct [cited 2019 Sep 21];95(6):437–42. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19747891>
9. Siegel MG. Editorial Commentary: “Defer No Time, Delays Have Dangerous Ends” (Henry VI,



- Shakespeare): Delayed Anterior Cruciate Ligament Reconstruction Has Consequences. *Arthroscopy - Journal of Arthroscopic and Related Surgery*. 2018;19:18–20.
10. Frobell RB, Roos HP, Roos EM, Roemer FW, Ranstam J, Lohmander LS. Treatment for acute anterior cruciate ligament tear: five year outcome of randomised trial. *BMJ* [Internet]. 2013 Jan 24 [cited 2018 Nov 12];346(jan24 1):f232–f232. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23349407>
  11. Karikis I, Åhlén M, Sernert N, Ejerhed L, Rostgård-Christensen L, Kartus J. The Long-Term Outcome After Early and Late Anterior Cruciate Ligament Reconstruction. *Arthrosc - J Arthrosc Relat Surg*. 2018;34(6):1907–17.
  12. Wellsandt E, Failla MJ, Axe MJ, Snyder-Mackler L. Does Anterior Cruciate Ligament Reconstruction Improve Functional and Radiographic Outcomes Over Nonoperative Management 5 Years After Injury? *Am J Sports Med*. 2018;46 (9):2103–12.
  13. van Yperen DT, Reijman M, van Es EM, Bierma-Zeinstra SMA, Meuffels DE. Twenty-Year Follow-up Study Comparing Operative Versus Nonoperative Treatment of Anterior Cruciate Ligament Ruptures in High-Level Athletes. *Am J Sports Med*. 2018;46(5):1129–36.
  14. Thompson SM, Salmon LJ, Waller A, Linklater J, Roe JP, Pinczewski LA. Twenty-Year Outcome of a Longitudinal Prospective Evaluation of Isolated Endoscopic Anterior Cruciate Ligament Reconstruction With Patellar Tendon or Hamstring Autograft. *Am J Sports Med* [Internet]. 2016 Dec 19 [cited 2019 Sep 21];44(12):3083–94. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27492972>
  15. Thompson S, Salmon L, Waller A, Linklater J, Roe J, Pinczewski L. Twenty-Year Outcomes of a Longitudinal Prospective Evaluation of Isolated Endoscopic Anterior Cruciate Ligament Reconstruction With Patellar Tendon Autografts. *Am J Sports Med* [Internet]. 2015 Sep 17 [cited 2019 Sep 21];43(9):2164–74. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26187130>
  16. Harris JD, Abrams GD, Bach BR, Williams D, Heidloff D, Bush-Joseph CA, et al. Return to Sport After ACL Reconstruction. *Orthopedics* [Internet]. 2014 Feb 1 [cited 2019 Sep 21];37(2):e103–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24679194>
  17. Filbay SR. Early ACL reconstruction is required to prevent additional knee injury: a misconception not supported by high-quality evidence. *Br J Sports Med* [Internet]. 2019 Apr 1 [cited 2019 Sep 21];53(8):459–61. Available from:

- <http://www.ncbi.nlm.nih.gov/pubmed/30377174>
18. Racine J, Aaron RK. Post-traumatic osteoarthritis after ACL injury. *R I Med J* (2013) [Internet]. 2014 Nov 3 [cited 2019 Sep 21];97(11):25–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25365816>
  19. Ajuied A, Wong F, Smith C, Norris M, Earnshaw P, Back D, et al. Anterior Cruciate Ligament Injury and Radiologic Progression of Knee Osteoarthritis. *Am J Sports Med* [Internet]. 2014 Sep 8 [cited 2019 Sep 21];42(9):2242–52. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24214929>
  20. Coughlin TR, Kennedy OD. The role of subchondral bone damage in post-traumatic osteoarthritis. *Ann N Y Acad Sci* [Internet]. 2016 Nov [cited 2019 Sep 21];1383(1):58–66. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27671712>
  21. Jungmann PM, Baum T, Nevitt MC, Nardo L, Gersing AS, Lane NE, et al. Degeneration in ACL Injured Knees with and without Reconstruction in Relation to Muscle Size and Fat Content—Data from the Osteoarthritis Initiative. Kellermayer MS, editor. *PLoS One* [Internet]. 2016 Dec 5 [cited 2019 Sep 21];11(12):e0166865. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27918596>
  22. Naraghi A WL. MRI evaluation of the postoperative knee: special considerations and pitfalls. *Clin Sport Med*. 2006;25(4):703-25.
  23. Farshad-Amacker NA PH. MRI of knee ligament injury and reconstruction. *J Magn Reson Imaging*. 2013;38(4):757–73.
  24. Bining J, Andrews G FB. The ABCs of the anterior cruciate ligament: a primer for magnetic resonance imaging assessment of the normal, injured and surgically repaired anterior cruciate ligament. *Br J Sport Med*. 2009;43(11):856–62.
  25. Van Dyck P, Vanhoenacker FM, Gielen JL, Dossche L, Van Gestel J, Wouters K, et al. Three tesla magnetic resonance imaging of the anterior cruciate ligament of the knee: can we differentiate complete from partial tears? *Skeletal Radiol* [Internet]. 2011 Jun 8 [cited 2019 Sep 21];40(6):701–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20931190>
  26. Naraghi A, White LM. MR Imaging of Cruciate Ligaments. *Magn Reson Imaging Clin N Am* [Internet]. 2014 Nov [cited 2019 Sep 21];22(4):557–80. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25442023>

27. Vlychou M, Hantes M, Michalitsis S, Tsezou A, Fezoulidis I V., Malizos K. Chronic anterior cruciate ligament tears and associated meniscal and traumatic cartilage lesions: evaluation with morphological sequences at 3.0 T. *Skeletal Radiol* [Internet]. 2011 Jun 27 [cited 2019 Sep 21];40(6):709–16. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20978758>
28. Casagrande BU, Maxwell NJ, Kavanagh EC, Towers JD, Shen W FF. Normal appearance and complications of double-bundle and selective-bundle anterior cruciate ligament reconstructions using optimal MRI techniques. *AJR Am J Roentgenol*. 2009;192(5):1407–15.
29. Eggerding V, Meuffels DE, van Es E, Van Arkel ER, van de Brand I, van Linge J, et al. Operative versus Nonoperative Treatment for Acute Anterior Cruciate Ligament Tears: A Multicenter Randomized Trial [Internet]. Dutch trial Register number: NTR 2746. 2019 [cited 2019 Sep 21]. Available from: <https://www.isakos.com/2019/Abstract/10949>
30. Frobell RB, Roos EM, Roos HP, Ranstam J, Lohmander LS. A randomized trial of treatment for acute anterior cruciate ligament tears. *N Engl J Med*. 2010;363(4):331–42.
31. Frobell RB, Roos HP, Roos EM, Roemer FW, Ranstam J, Lohmander LS. Treatment for acute anterior cruciate ligament tear: Five year outcome of randomised trial. *Br J Sports Med*. 2015;49(10):700.
32. Smith TO, Postle K, Penny F, McNamara I, Mann CJV. Is reconstruction the best management strategy for anterior cruciate ligament rupture? A systematic review and meta-analysis comparing anterior cruciate ligament reconstruction versus non-operative treatment. *Knee* [Internet]. 2014 Mar [cited 2019 Sep 21];21(2):462–70. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24238648>
33. Sanders TL, Kremers HM, Bryan AJ, Fruth KM, Larson DR, Pareek A, et al. Is Anterior Cruciate Ligament Reconstruction Effective in Preventing Secondary Meniscal Tears and Osteoarthritis? *Am J Sports Med*. 2016;44(7):1699–707.
34. NCT02980367. ACL SNNAP Trial: ACL Surgery Necessity in Non Acute Patients. <https://clinicaltrials.gov/show/nct02980367>. 2016;
35. O’Shea JJ, Shelbourne KD. Repair of Locked Bucket-Handle Meniscal Tears in Knees with Chronic Anterior Cruciate Ligament Deficiency. *Am J Sports Med* [Internet]. 2003 Mar 30 [cited 2019 Sep 21];31(2):216–20. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12642255>
36. Wierer G, Runer A, Hoser C, Herbst E, Gföller P, Fink C. Acute ACL reconstruction in patients

- over 40 years of age. *Knee Surgery, Sport Traumatol Arthrosc* [Internet]. 2017 May 24 [cited 2019 Sep 21];25(5):1528–34. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27778041>
37. Cinque ME, Chahla J, Moatshe G, DePhillipo NN, Kennedy NI, Godin JA, et al. Outcomes and Complication Rates After Primary Anterior Cruciate Ligament Reconstruction Are Similar in Younger and Older Patients. *Orthop J Sport Med* [Internet]. 2017 Oct 2 [cited 2019 Sep 21];5(10):232596711772965. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29051896>
  38. Olsson O, Isacsson A, Englund M, Frobell RB. Epidemiology of intra- and peri-articular structural injuries in traumatic knee joint hemarthrosis – data from 1145 consecutive knees with subacute MRI. *Osteoarthr Cartil*. 2016 Nov 1;24(11):1890–7.
  39. Raheem O, Philpott J, Ryan W, O’Brien M. Anatomical variations in the anatomy of the posterolateral corner of the knee. *Knee Surgery, Sport Traumatol Arthrosc* [Internet]. 2007 Jul 19 [cited 2019 Sep 26];15(7):895–900. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17641923>
  40. Stannard JP. Evaluation and treatment of medial instability of the knee. *Sport Med Arthrosc*. 2015;23(2):91–5.
  41. Svantesson E, Hamrin Senorski E, Alentorn-Geli E, Westin O, Sundemo D, Grassi A, et al. Increased risk of ACL revision with non-surgical treatment of a concomitant medial collateral ligament injury: a study on 19,457 patients from the Swedish National Knee Ligament Registry. *Knee Surgery, Sport Traumatol Arthrosc*. 2019 Aug 1;27(8):2450–9.
  42. Schmitz MA1, Rouse LM Jr DK. The management of meniscal tears in the ACL-deficient knee. *Clin Sport Med*. 1996;15(3):573–93.
  43. Balasingam S, Sernert N, Magnusson H, Kartus J. Patients With Concomitant Intra-articular Lesions at Index Surgery Deteriorate in Their Knee Injury and Osteoarthritis Outcome Score in the Long Term More Than Patients With Isolated Anterior Cruciate Ligament Rupture: A Study From the Swedish National Anterior Cruciate Ligament Register. *Arthrosc - J Arthrosc Relat Surg*. 2018 May 1;34(5):1520–9.
  44. Fischer SP, Fox JM, Del PW, Friedman MJ, Snyder SJ FR. Accuracy of diagnoses from magnetic resonance imaging of the knee. A multi-center analysis of one thousand and fourteen patients. *J Bone Jt Surg Am*. 1991;73:2–10.

45. Jah AAE, Keyhani S, Zarei R MA. Accuracy of MRI in comparison with clinical and arthroscopic findings in ligamentous and meniscal injuries of the knee. *Acta Orthop Belg.* 2005;71:189–96.
46. Rayan F, Bhonsle S SD. Clinical, MRI, and arthroscopic correlation in meniscal and anterior cruciate ligament injuries. *Int Orthop.* 2009;33:129–32.
47. Shelbourne KD, Porter DA, Clingman JA, McCarroll JR RA. Low-velocity knee dislocation. *Orthop Rev.* 1991;20(11):995–1004.
48. Azar FM, Brandt JC, Miller Iii RH, Phillips BB. Ultra-Low-Velocity Knee Dislocations. *Am J Sport Med* [Internet]. 2011 [cited 2018 Jul 1];30(10):2170–4. Available from: <http://journals.sagepub.com/doi/pdf/10.1177/0363546511414855>
49. Werner BC, Gwathmey FW, Higgins ST, Hart JM, Miller MD. Ultra-Low Velocity Knee Dislocations. *Am J Sports Med* [Internet]. 2014 Feb 8 [cited 2018 Jul 22];42(2):358–63. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24214926>
50. Carr JB, Werner BC, Miller MD, Gwathmey FW. Knee Dislocation in the Morbidly Obese Patient. *J Knee Surg.* 2016 May 1;29(4):278–86.
51. Frassica FJ, Sim FH, Staeheli JW, Pairolo PC. Dislocation of the knee. *Clin Orthop Relat Res* [Internet]. 1991 Feb [cited 2018 Jul 23];263:200–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/1993376>
52. Medina O, Arom GA, Yerosian MG, Petrigliano FA, McAllister DR. Vascular and Nerve Injury After Knee Dislocation: A Systematic Review. *Clin Orthop Relat Res* [Internet]. 2014 Sep 20 [cited 2019 Sep 26];472(9):2621–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24554457>
53. Johnson JP, Kleiner J, Klinge SA, McClure PK, Hayda RA, Born CT. Increased Incidence of Vascular Injury in Obese Patients With Knee Dislocations. In: *Journal of Orthopaedic Trauma.* Lippincott Williams and Wilkins; 2018. p. 82–7.
54. Medina Bs O, Arom Bs GA, Yerosian MG, Petrigliano FA, Mcallister DR. Vascular and Nerve Injury After Knee Dislocation A Systematic Review. *Clin Orthop Relat Res* [Internet]. 2014 [cited 2018 Jul 1];472(9):2621–9. Available from: [https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4117866/pdf/11999\\_2014\\_Article\\_3511.pdf](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4117866/pdf/11999_2014_Article_3511.pdf)
55. Howells NR, Brunton LR, Robinson J, Porteous AJ, Eldridge JD MJ. Acute knee dislocation: an evidence based approach to the management of the multiligament injured knee. *Injury.*

- 2011;42(11):1198–204.
56. Anand S. Fracture Dislocations of the Knee. In: *The Multiple Ligament Injured Knee* [Internet]. Cham: Springer International Publishing; 2019 [cited 2019 Sep 26]. p. 395–412. Available from: [http://link.springer.com/10.1007/978-3-030-05396-3\\_28](http://link.springer.com/10.1007/978-3-030-05396-3_28)
  57. Reider B. Medial Collateral Ligament Injuries in Athletes. *Sport Med*. 1996;21:147–56.
  58. Noyes, F.R. and Barber-Westin SD. The Treatment of Acute Combined Ruptures of the Anterior Cruciate and Medial Ligaments of the Knee. *Am J Sport Med*. 1995;23:380–91.
  59. Kakarlapudi, T.K. and Bickerstaff DR. Knee Instability: isolated and complex. *WestJMed*. 2001;174:266–72.
  60. Robinson JR, Bull AM, Thomas RR AA. The role of the medial collateral ligament and posteromedial capsule in controlling knee laxity. *Am J Sport Med*. 2006;34(11):1815–23.
  61. Jari, S. and Shelbourne KD. Non operative or delayed surgical treatment of combined cruciate ligaments and medial side knee injuries. *Sport Med Arthrosc Rev* 9. 2001;185–92.
  62. Zaffagnini S, Bonanzinga T, Marcheggiani Muccioli GM, Giordano G, Bruni D, Bignozzi S, et al. Does chronic medial collateral ligament laxity influence the outcome of anterior cruciate ligament reconstruction?: a prospective evaluation with a minimum three-year follow-up. *J Bone Joint Surg Br* [Internet]. 2011 Aug [cited 2019 Dec 30];93(8):1060–4. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21768629>
  63. Kannus P. Long-term results of conservatively treated medial collateral ligament injuries of the knee joint. *Clin Orthop*. 1988(226):103–112.
  64. Fetto, J.F. and Marshall JL. Medial collateral ligament injuries of the knee: a rationale for treatment. *ClinOrthop*. 1978;206–18.
  65. Arthur A, , LaPrade RF AJ. Proximal Tibial Opening Wedge Osteotomy as the Initial Treatment for Chronic Posterolateral Corner Deficiency in the Varus Knee A Prospective Clinical Study. *Am J Sport Med*. 2007;35(11).
  66. Wilde J, Bedi, A, Alcheck D. Revision Anterior Cruciate Ligament Reconstruction. *Sport Heal*. 2014;6(6):504–518.
  67. Moorman CT 3rd LR. Anatomy and biomechanics of the posterolateral corner of the knee. *J Knee Surg*. 2005;18:137–145.

68. LaPrade RF, Resig S, Wentorf F LJ. The effects of grade III posterolateral knee complex injuries on anterior cruciate ligament graft force: a biomechanical analysis. *Am J Sport Med.* 1999;27:469.
69. Chen FS, Rokito AS PM. Acute and chronic posterolateral rotatory instability of the knee. *J Am Acad Orthop Surg.* 2000;8:97–110.
70. Chahla J, Murray IR, Robinson J, Lagae K, Margheritini F, Fritsch B, et al. Posterolateral corner of the knee: an expert consensus statement on diagnosis, classification, treatment, and rehabilitation. *Knee Surgery, Sport Traumatol Arthrosc.* 2019 Aug 1;27(8):2520–9.
71. Covey DC. Injuries of the posterolateral corner of the knee. *J Bone Jt Surg Am.* 2001;83:106–118.
72. Levy BA, Dajani KA, Morgan JA, Shah JP, Dahm DiL, Stuart MJ. Repair Versus Reconstruction of the Fibular Collateral Ligament and Posterolateral Corner in the Multiligament-Injured Knee. In: *American Journal of Sports Medicine.* SAGE Publications Inc.; 2010. p. 804–9.
73. Stannard JP, Brown SL, Farris RC, McGwin G, Volgas DA. The posterolateral corner of the knee: Repair versus reconstruction. *Am J Sports Med.* 2005 Jun;33(6):881–8.
74. Westermann RW, Marx RG, Spindler KP, Huston LJ, Amendola A, Andrish JT, et al. No Difference Between Posterolateral Corner Repair and Reconstruction With Concurrent ACL Surgery: Results From a Prospective Multicenter Cohort. *Orthop J Sport Med.* 2019 Jul 1;7(7).
75. Dedmond BT AL. Operative versus nonoperative treatment of knee dislocations: a meta-analysis. *Am J Knee Surg.* 2001;1:33–8.
76. Panigrahi R, Kumari Mahapatra A, Priyadarshi A, Singha Das D, Palo N RBM. Outcome of Simultaneous Arthroscopic Anterior Cruciate Ligament and Posterior Cruciate Ligament Reconstruction With Hamstring Tendon Autograft: A Multicenter Prospective Study. *Asian J Sport Med.* 2016;7(1):e29287.
77. Noyes FR B-WS. Reconstruction of the anterior and posterior cruciate ligaments after knee dislocation. Use of early protected postoperative motion to decrease arthrofibrosis. *Am J Sport Med.* 1997;2:769–78.
78. Levy BA, Dajani KA, Whelan DB et al. Decision making in the multi-ligamentinjured knee: an evidence-based systematic review. *Arthroscopy.* 2009;2:430–8.

79. Spindler KP, Schils JP, Bergfeld JA et al. Prospective study of osseous, articular, and meniscal lesions in recent anterior cruciate ligament tears by magnetic resonance imaging and arthroscopy. *Am J Sport Med.* 1993;21:551–7.
80. Bellabarba C, Bush-Joseph CA BBJ. Patterns of meniscal injury in the anterior cruciate-deficient knee: a review of the literature. *Am J Orthop (Belle Mead NJ).* 1997;26:18–23.
81. Shoemaker SC MK. The role of the meniscus in the anterior-posterior stability of the loaded anterior cruciate-deficient knee. Effects of partial versus total excision. *J Bone Jt Surg Am.* 1986;68(1):71–9.
82. Sisto DJ, Warren RF. Complete knee dislocation. A follow-up study of operative treatment. *Clin Orthop Relat Res [Internet].* 1985 Sep [cited 2018 Jul 23];198:94–101. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/4028570>
83. Chhadia AM, Inacio MCS, Maletis GB et al. Are meniscus and cartilage injuries related to time to anterior cruciate ligament reconstruction? *Am J Sport Med.* 2011;39:1894–1899.
84. Hagino T, Ochiai S, Senga S, Yamashita T, Wako M, Ando T HH. Meniscal tears associated with anterior cruciate ligament injury. *Arch Orthop Trauma Surg.* 2015;(135(12)):1701–6.
85. Walter RP, Dhadwal AS, Schranz P M V. The outcome of all-inside meniscal repair with relation to previous anterior cruciate ligament reconstruction. *Knee.* 2014;21(6):1156–9.
86. Tanaka M, Vyas D, Moloney G, Bedi A, Pearle AD M V. What does it take to have a high-grade pivot shift? *Knee Surg Sport Traumatol Arthrosc.* 2012;20(4):737–42.
87. Song GY1, Zhang H1, Wang QQ2, Zhang J1, Li Y1 FH. Risk Factors Associated With Grade 3 Pivot Shift After Acute Anterior Cruciate Ligament Injuries. *Am J Sport Med.* 2016;44(2):362–9.
88. Musahl V, Citak M, O’Loughlin PF, Choi D, Bedi A PA. The effect of medial versus lateral meniscectomy on the stability of the anterior cruciate ligament-deficient knee. *Am J Sport Med.* 2010;38(8):1591–7.
89. Shelbourne KD DM. Comparison of partial meniscectomy versus meniscus repair for bucket-handle lateral meniscus tears in anterior cruciate ligament reconstructed knees. *Arthroscopy.* 2004;20:581–5.
90. Wu H, Hackett T RJ. Effects of meniscal and articular surface status on knee stability, function, and symptoms after anterior cruciate ligament reconstruction: A long-term prospective study.



- Am J Sport Med. 2002;30:845–50.
91. Aglietti P, Zaccherotti G, De BP TI. A comparison between medial meniscus repair, partial meniscectomy, and normal meniscus in anterior cruciate ligament reconstructed knees. *Clin Orthop Relat Res.* 1994;165–73.
  92. Kennedy MI, LaPrade RF, Marwan by. Editorial Commentary: Is the Anterolateral Ligament Always Injured in Knee Dislocations? Does It Matter? *Arthrosc J Arthrosc Relat Surg* [Internet]. 2018 [cited 2018 Jul 7];34(6):1898–9. Available from: <https://doi.org/10.1016/j.arthro.2018.04.004>
  93. Orfaly R, McConkey J RW. The fate of meniscal tears after anterior cruciate ligament reconstruction. *Clin J Sport Med.* 1998;8:102–105.
  94. Biant LC, McNicholas MJ, Sprowson AP ST. The surgical management of symptomatic articular cartilage defects of the knee: Consensus statements from United Kingdom knee surgeons. *Knee.* 2015;22(5):446–9.
  95. Mall NA, Chalmers PN, Moric M, Tanaka MJ, Cole BJ, Bach BR PG. Incidence and trends of anterior cruciate ligament reconstruction in the United States. *Am J Sport Med.* 2014;42:2362–70.
  96. Dodwell ER, LaMont LE, Green DW, Pan TJ, Marx RG LS. 20 years of pediatric anterior cruciate ligament reconstruction in New York State. *Am J Sport Med.* 2014;42:675–80.
  97. Werner BC, Yang S LA. Trends in pediatric and adolescent anterior cruciate injury and reconstruction. *J Pediatr Orthop.* 2016;36:447–52.
  98. Nogaro MC, Abram SGF, Alvand A, Bottomley N, Jackson WFM, Price A. Paediatric and adolescent anterior cruciate ligament reconstruction surgery. *Bone Joint J* [Internet]. 2020 Feb 1 [cited 2020 Feb 18];102-B(2):239–45. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/32009437>
  99. Gilchrist J, Mandlebaum BR, Melancon H, Ryan GW, Silvers HJ, Griffin LY, Watanabe DS, Dick RW DJ. A randomized controlled trial to prevent noncontact anterior cruciate ligament injury in female collegiate soccer players. *Am J Sport Med.* 2008;36:1476–83.
  100. Wingfield K. Neuromuscular training to prevent knee injuries in adolescent female soccer players. *Clin J Sport Med.* 2013;23:407–8.

101. Stevenson JH, Beattie CS, Schwartz JB BB. Assessing the effectiveness of neuromuscular training programs in reducing the incidence of anterior cruciate ligament injuries in female athletes: A systematic review. *Am J Sport Med.* 2015;43:482–90.
102. Fabricant PD, Jones KJ, Delos D, Cordasco FA, Maex RG, Pearle AD, Warren RF GD. Reconstruction of the anterior cruciate ligament in the skeletally immature athlete: A review of current concepts: AAOS exhibit selection. *J Bone Jt Surg Am.* 2013;95:e28(1-13).
103. Ramski DE, Kanj WW, Franklin CC, Baldwin KD GT. Anterior cruciate ligament tears in children and adolescents: A meta-analysis of nonoperative versus operative treatment. *Am J Sport Med.* 2014;42:2769–76.
104. Henry J, Chotel F, Chouteau J, Fessy MH, Berard J MB. Rupture of the anterior cruciate ligament in children: early reconstruction with open physes or delayed reconstruction to skeletal maturity? *Knee Surg Sport Traumatol Arthrosc.* 2009;17:748–55.
105. Lawrence JTR, Argawal N GT. Degeneration of the knee joint in skeletally immature patients with a diagnosis of an anterior cruciate ligament tear: Is there harm in delay of treatment? *Am J Sport Med.* 2011;39:2582–7.
106. Anderson AF AC. Correlation of meniscal and articular cartilage injuries in children and adolescents with timing of anterior cruciate ligament reconstruction. *Am J Sport Med.* 2015;43:275–81.
107. Smith JO, Yasen S, Palmer HC, Lord BR, Britton E WA. Paediatric ACL repair reinforced with temporary internal bracing. *Knee Surg Sport Traumatol Arthrosc.* 2016;24:1845–51.
108. Kocher MS, Smith JT, Zoric BJ, Lee B ML. Transphyseal anterior cruciate ligament reconstruction in skeletally immature prepubescent adolescents. *J Bone Jt Surg Am.* 2007;89:2632–9.
109. Liddle AD, Imbuldeniya AM HD. Transphyseal reconstruction of the anterior cruciate ligament in prepubescent children. *J Bone Jt Surg.* 2008;90:1317–22.
110. Courvoisier A GM. Good surgical outcome of transphyseal ACL reconstruction in skeletally immature patients using four strand hamstring graft. *Knee Surg Sport Traumatol Arthrosc.* 2011;19:588–91.
111. Nikolaou P, Kalliakmanis A, Bousgas D ZS. Intraarticular stabilization following anterior cruciate ligament injury in children and adolescents. *Knee Surg Sport Traumatol Arthrosc.* 2011;19:801–5.

112. Hui C, Roe J, Ferguson D, Waller A, Salmon L PL. Outcome of anatomic transphyseal anterior cruciate ligament reconstruction in Tanner stage 1 and 2 patients with open physes. *Am J Sport Med.* 2012;40:1093–8.
113. Calvo R, Figueroa D, Gili F, Vaisman A, Mococain P, Espinosa M, Leon A AS. Transphyseal anterior cruciate ligament reconstruction in patients with open physes: 10 year follow up study. *Am J Sport Med.* 2015;43:289–94.
114. Morgan MD, Salmon LJ, Waller A, Roe JP PL. Fifteen year survival of endoscopic anterior cruciate ligament reconstruction in patients aged 18 years and younger. *Am J Sport Med.* 2016;44:384–92.
115. Kocher MS, Garg S ML. Physeal sparing reconstruction of the anterior cruciate ligament in skeletally immature prepubescent children and adolescents. *J Bone Jt Surg Am.* 2005;87:2371–9.
116. Willimon SC, Jones CR, Herzog MM, May, KH, Leake M BM. Micheli anterior cruciate ligament reconstruction in skeletally immature youths: A retrospective case series with a mean 3-year follow up. *Am J Sport Med.* 2015;43:2974–81.
117. Kocher MS, Heyworth BE, Fabricant PD, Tepolt FA, Micheli LJ. Outcomes of Physeal-Sparing ACL Reconstruction with Iliotibial Band Autograft in Skeletally Immature Prepubescent Children. *J Bone Joint Surg Am* [Internet]. 2018 Jul 5 [cited 2020 Jan 24];100(13):1087–94. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29975275>
118. Guzzanti V, Falciglia F SS. Physeal sparing intraarticular anterior cruciate ligament reconstruction in preadolescents. *Am J Sport Med.* 2003;31:949–53.
119. Anderson AF. Transepiphyseal replacement of the anterior cruciate ligament in skeletally immature patients. A preliminary report. *J Bone Jt Surg Am.* 2003;85:1255–63.
120. Lawrence JT, Bowers AL, Belding, J, Cody SR GT. All epiphyseal anterior cruciate ligament reconstruction in skeletally immature patients. *Clin Orthop Relat Res.* 2010;468:1971–7.
121. Demange MK & Camanh GL. Nonanatomic anterior cruciate ligament reconstruction with double stranded semitendinosus grafts in children wuth open physes; Minimum 15 year follow up. *Am J Sport Med.* 2014;42:2926–32.
122. Pierce TP, Issa K, Festa A, Scillia AJ MV. Pediatric anterior cruciate ligament reconstruction: A systematic review of transphyseal versus physeal sparing techniques. *Am J Sport Med.*

- 2017;45:488–494.
123. Moksnes H, Engebretsen L SR. The ESSKA paediatric anterior cruciate monitoring initiative. *Knee Surg Sport Traumatol Arthrosc.* 2016;24:680–7.
  124. Edwards TB, Greene CC, Baratta RV, Zieske A WR. The effect of placing a tensioned graft across open growth plates: A gross and histological analysis. *J Bone Jt Surg Am.* 2001;83:725–34.
  125. Chudik S, Beasley L, Potter H, Wickiewicz, Warren R RS. The influence of femoral technique for graft placement on anterior cruciate ligament reconstruction using a skeletally immature canine model with a rapidly growing physis. *Arthroscopy.* 2007;23:1309–19.
  126. Yoo WJ, Kocher MS ML. Growth plate disturbance after transphyseal reconstruction of the anterior cruciate ligament in skeletally immature adolescent patients: An MRI imaging study. *Orthop, J Pediatr.* 2011;32:691–6.
  127. Kocher MS, Saxon HS HW. Management and complications of anterior cruciate ligament injuries in skeletally immature patients: Survey of the Herodicus society and the ACL study group. *J Pediatr Orthop.* 2002;22:452–7.
  128. Chotel F, Henry J, Seil R, Chouteau J, Moyen B BJ. Growth disturbance without growth arrest after ACL reconstruction in children. *Knee Surg Sport Traumatol Arthrosc.* 2010;18:1496–500.
  129. Lawrence JT, West RL GW. Growth disturbance following ACL reconstruction with use of an epiphyseal femoral tunnel: A case report. *J Bone Jt Surg Am.* 2011;93:e39(1-6).
  130. Tomatsuri M, Yoshiya S, Kuroda R, Matsushita T KM. Limb deformity caused by distal femoral and proximal tibial growth arrest after ACL reconstruction in a child. *J Bone Jt Surg Case Connect.* 2015;5:e84.
  131. Shifflett GD, Green DW WR. Growth arrest following ACL reconstruction with hamstring autograft in skeletally immature patients: A review of 4 cases. *J Pediatr Orthop.* 2016;36:355–61.
  132. Volpi P, Cervellin M, Bait C, Prospero E, Mousa H, Redaelli A, Quaglia A DM. Transphyseal anterior cruciate ligament reconstruction in adolescents. *Knee Surg Sport Traumatol Arthrosc.* 2016;24:707–11.
  133. Kaeding CC, Pedroza AD, Reinke EK, Huston LJ M consortium & SK. Risk factors and predictors of subsequent ACL injury in either knee after ACL reconstruction: Prospective analysis of 2488

- primary ACL reconstructions from the MOON cohort. *Am J Sport Med.* 2015;43:1583–90.
134. Shelbourne KD, Gray T HM. Incidence of subsequent injury to either knee within 5 years after anterior cruciate ligament reconstruction with patellar tendon autograft. *Am J Sport Med.* 2009;37:246–51.
  135. Webster KE, Feller JA, Leigh WB RA. Younger patients are at increased risk for graft rupture and contralateral injury after anterior cruciate ligament reconstruction. *Am J Sport Med.* 2014;42:641.
  136. Andernord D, Desai N, Bjornsson H, Ylander M, Karlsson J SK. Patient predictors of early revision surgery after anterior cruciate ligament reconstruction: A cohort study of 16,930 patients with 2 year follow up. *Am J Sport Med.* 2015;43:121–7.
  137. Andernord D, Desai N, Bjornsson, Gillen S, Karlsson J SK. Predictors of contralateral anterior cruciate ligament reconstruction: A cohort study of 9,061 patients with 5 year follow up. *Am J Sport Med.* 2015;43:295–302.
  138. Wiggins ME, Fadale P, Barrach H, Ehrlich M, WW. Risk of secondary injury in younger athletes after anterior cruciate ligament reconstruction: a systematic review and meta-analysis. *Am J Sport Med.* 2016;44:1861–76.
  139. Rahr-Wagner L, Thillemann TM, Pedersen AB LM. Comparison of hamstring tendon and patellar tendon grafts in anterior cruciate ligament reconstruction in a nationwide population based cohort study: Results from the Danish registry of knee ligament reconstruction. *Am J Sport Med.* 2014;42:278–84.
  140. Persson A, Fjeldsgaard K, Gjertsen JE, Kjellsen A, Engebretsen L, Hole RM FJ. Increased risk of revision with hamstring tendon grafts compared with patellar tendon grafts after anterior cruciate ligament reconstruction: A study of 12,643 patients from the Norwegian cruciate ligament registry. *Am J Sport Med.* 2014;42:285–91.
  141. Ho B, Edmonds EW, Chambers HG, Bastrom T PA. Risk factors for early ACL reconstruction failure in pediatric and adolescent patients: A review of 561 cases. *J Paedr Orthop.* 2018;38(7):388–92.
  142. Maletis GB, Chen J, Inacio MCS FT. Age related risk factors for revision anterior cruciate ligament reconstruction: A cohort study of 21,304 patients from the Kaiser Permanente anterior cruciate ligament registry. *Am J Sport Med.* 2016;44:331–6.

143. Goddard M, Bowman N, Salmon LJ, Waller A, Roe J P LA. Endoscopic anterior cruciate ligament reconstruction in children using living donor hamstring tendon allografts. *Am J Sport Med.* 2013;41:567.
144. Hewett TE, Di Stasi SL MG. Current concepts for injury prevention in athletes after anterior cruciate ligament reconstruction. *Am J Sport Med.* 2013;41:216–24.
145. Herd H. The Benefits of a Preoperative MRSA Swabbing Protocol. *J Perianesthesia Nurs.* 2017 Apr 1;32(2):134–9.
146. Adams D, Logerstedt D, Hunter-Giordano A, Axe M S-MLC concepts for anterior cruciate ligament reconstruction: A criterion based rehabilitation progression J of O and 42; Current concepts for anterior cruciate ligament reconstruction: A criterion based rehabilitation progression. *J Orthop Sport Phys Ther.* 2012;42:601–14.
147. Quelard B, Sonnery-Cottet B, Zayni R, Ogassawara R, Prost T, Chambat P. Preoperative factors correlating with prolonged range of motion deficit after anterior cruciate ligament reconstruction. *Am J Sports Med.* 2010 Oct;38(10):2034–9.
148. Failla MJ, Logerstedt DS, Grindem H, Axe MJ, Risberg MA, Engebretsen L, et al. Does Extended Preoperative Rehabilitation Influence Outcomes 2 Years after ACL Reconstruction? A Comparative Effectiveness Study between the MOON and Delaware-Oslo ACL Cohorts: *Am J Sports Med.* 2016;44(10):2608–14.
149. Consent: Supported Decision-Making — Royal College of Surgeons [Internet]. [cited 2020 Jan 5]. Available from: <https://www.rcseng.ac.uk/standards-and-research/standards-and-guidance/good-practice-guides/consent/>
150. Monk AP, Davies LJ, Hopewell S, Harris K, Beard DJ, Price AJ. Surgical versus conservative interventions for treating anterior cruciate ligament injuries. *Cochrane Database Syst Rev* [Internet]. 2016 Apr 3 [cited 2019 Sep 21];4:CD011166. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27039329>
151. Svantesson E, Hamrin Senorski E, Östergaard M, Grassi A, Krupic F, Westin O, et al. Graft Choice for Anterior Cruciate Ligament Reconstruction With a Concomitant Non-surgically Treated Medial Collateral Ligament Injury Does Not Influence the Risk of Revision. *Arthrosc - J Arthrosc Relat Surg.* 2020 Jan 1;36(1):199–211.
152. Grassi A, Nitri M, Moulton SG, Marcheggiani GM, Muccioli, Bondi A, et al. Does the type of graft

- affect the outcome of revision anterior cruciate ligament reconstruction? a meta-analysis of 32 studies. Vol. 99B, *Bone and Joint Journal*. British Editorial Society of Bone and Joint Surgery; 2017. p. 714–23.
153. Wang H De, Zhang H, Wang TR, Zhang WF, Wang FS, Zhang YZ. Comparison of clinical outcomes after anterior cruciate ligament reconstruction with hamstring tendon autograft versus soft-tissue allograft: A meta-analysis of randomised controlled trials. Vol. 56, *International Journal of Surgery*. Elsevier Ltd; 2018. p. 174–83.
  154. Samitier G, Marcano AI, Alentorn-Geli E, Cugat R, Farmer KW, Moser MW. Failure of anterior cruciate ligament reconstruction. Vol. 3, *Archives of Bone and Joint Surgery*. Mashhad University of Medical Sciences; 2015. p. 220–40.
  155. Gaskill T, Pullen M, Bryant B, Sicignano N, Evans AM, DeMaio M. The Prevalence of Symptomatic Deep Venous Thrombosis and Pulmonary Embolism After Anterior Cruciate Ligament Reconstruction. *Am J Sports Med* [Internet]. 2015 Nov [cited 2020 Jan 5];43(11):2714–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26391861>
  156. Jameson SS, Downen D, James P, Serrano-Pedraza I, Reed MR, Deehan D. Complications following anterior cruciate ligament reconstruction in the English NHS. *Knee* [Internet]. 2012 Jan [cited 2018 Nov 12];19(1):14–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21216599>
  157. Jameson SS, Downen D, James P, Serrano-Pedraza I, Serrano-Pedraza D, Reed MR, et al. The burden of arthroscopy of the knee: a contemporary analysis of data from the English NHS.[Erratum appears in *J Bone Joint Surg Br*. 2011 Dec;93(12):1679 Note: Serrano-Pedraza, Deehan [corrected to Serrano-Pedraza, I]]. *J Bone Jt Surg - Br Vol* [Internet]. 2011 [cited 2020 Jan 5];93(10):1327–33. Available from: <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&D=med5&AN=21969430>
  158. Wiggins AJ, Grandhi RK, Schneider DK, Stanfield D, Webster KE, Myer GD. Risk of Secondary Injury in Younger Athletes after Anterior Cruciate Ligament Reconstruction. Vol. 44, *American Journal of Sports Medicine*. SAGE Publications Inc.; 2016. p. 1861–76.
  159. Andernord D, Desai N, Björnsson H, Gillén S, Karlsson J, Samuelsson K. Predictors of contralateral anterior cruciate ligament reconstruction: A cohort study of 9061 patients with 5-year follow-up. *Am J Sports Med*. 2015 Feb 4;43(2):295–302.

160. Grindem H, Snyder-Mackler L, Moksnes H, Engebretsen L, Risberg MA. Simple decision rules can reduce reinjury risk by 84% after ACL reconstruction: The Delaware-Oslo ACL cohort study. *Br J Sports Med.* 2016;
161. Van Melick N, Van Cingel REH, Brooijmans F, Neeter C, Van Tienen T, Hullegie W, et al. Evidence-based clinical practice update: Practice guidelines for anterior cruciate ligament rehabilitation based on a systematic review and multidisciplinary consensus. Vol. 50, *British Journal of Sports Medicine.* BMJ Publishing Group; 2016. p. 1506–15.
162. Davies GJ, McCarty E, Provencher M, Manske RC. ACL Return to Sport Guidelines and Criteria. Vol. 10, *Current Reviews in Musculoskeletal Medicine.* Humana Press Inc.; 2017. p. 307–14.
163. Pasquarella C, Barchitta M, D’Alessandro D, Cristina ML, Mura I, Nobile M, et al. Heating, ventilation and air conditioning (HVAC) system, microbial air contamination and surgical site infection in hip and knee arthroplasties: The GISIO-SItI Ischia study. *Ann di Ig.* 2018;30(5):22–35.
164. Bischoff P, Kubilay NZ, Allegranzi B, Egger M, Gastmeier P. Effect of laminar airflow ventilation on surgical site infections: a systematic review and meta-analysis. *Lancet Infect Dis.* 2017 May 1;17(5):553–61.
165. McHugh SM, Hill ADK, Humphreys H. Laminar airflow and the prevention of surgical site infection. More harm than good? Vol. 13, *Surgeon.* Elsevier Ltd; 2015. p. 52–8.
166. Chapter 6: Guidelines for the Provision of Anaesthesia Services for Day Surgery 2019 | The Royal College of Anaesthetists [Internet]. [cited 2020 Jan 5]. Available from: <https://www.rcoa.ac.uk/gpas/chapter-6>
167. Abdallah FW, Brull R, Joshi GP. Pain Management for Ambulatory Arthroscopic Anterior Cruciate Ligament Reconstruction: Evidence-Based Recommendations From the Society for Ambulatory Anesthesia. *Anesth Analg.* 2019 Apr 1;128(4):631–40.
168. Smith I, Kranke P, Murat I, Smith A, O’Sullivan G, Søreide E, et al. Perioperative fasting in adults and children: Guidelines from the european society of anaesthesiology. *Eur J Anaesthesiol.* 2011;28(8):556–69.
169. Joshi GP, Kehlet H, Beloeil H, Bonnet F, Fischer B, Hill A, et al. Guidelines for perioperative pain management: Need for re-evaluation. Vol. 119, *British Journal of Anaesthesia.* Elsevier Ltd; 2017. p. 720–2.



170. Mardani-Kivi M, Mobarakeh MK, Keyhani S, Motlagh KH, Ekhtiari KS. Is gabapentin effective on pain management after arthroscopic anterior cruciate ligament reconstruction? A triple blinded randomized controlled trial. *Arch Bone Jt Surg*. 2013 Sep 1;1(1):18–22.
171. Secrist ES, Freedman KB, Ciccotti MG, Mazur DW, Hammoud S. Pain management after outpatient anterior cruciate ligament reconstruction: A systematic review of randomized controlled trials. Vol. 44, *American Journal of Sports Medicine*. SAGE Publications Inc.; 2016. p. 2435–47.
172. Swank KR, DiBartola AC, Everhart JS, Kaeding CC, Magnussen RA, Flanigan DC. The Effect of Femoral Nerve Block on Quadriceps Strength in Anterior Cruciate Ligament Reconstruction: A Systematic Review. Vol. 33, *Arthroscopy - Journal of Arthroscopic and Related Surgery*. W.B. Saunders; 2017. p. 1082-1091.e1.
173. Luo TD, Ashraf A, Dahm DL, Stuart MJ, McIntosh AL. Femoral nerve block is associated with persistent strength deficits at 6 months after anterior cruciate ligament reconstruction in pediatric and adolescent patients. *Am J Sports Med*. 2015 Feb 4;43(2):331–6.
174. Vorobeichik L, Brull R, Joshi GP, Abdallah FW. Evidence Basis for Regional Anesthesia in Ambulatory Anterior Cruciate Ligament Reconstruction. *Anesth Analg* [Internet]. 2019 Jan [cited 2020 Jan 11];128(1):58–65. Available from: <http://insights.ovid.com/crossref?an=00000539-201901000-00013>
175. Lynch JR, Okoroha KR, Lizzio V, Yu CC, Jildeh TR, Moutzouros V. Adductor Canal Block Versus Femoral Nerve Block for Pain Control After Anterior Cruciate Ligament Reconstruction: A Prospective Randomized Trial. *Am J Sports Med*. 2019 Feb 1;47(2):355–63.
176. Bailey L, Griffin J, Elliott M, Wu J, Papavasiliou T, Harner C, et al. Adductor Canal Nerve Versus Femoral Nerve Blockade for Pain Control and Quadriceps Function Following Anterior Cruciate Ligament Reconstruction With Patellar Tendon Autograft: A Prospective Randomized Trial. *Arthrosc - J Arthrosc Relat Surg*. 2019 Mar 1;35(3):921–9.
177. Ahl MS El. Femoral nerve block versus adductor canal block for postoperative pain control after anterior cruciate ligament reconstruction: A randomized controlled double blind study. *Saudi J Anaesth*. 2015;9(3):279–82.
178. Runner RP, Boden SA, Godfrey WS, Premkumar A, Samady H, Gottschalk MB, et al. Quadriceps Strength Deficits After a Femoral Nerve Block Versus Adductor Canal Block for Anterior Cruciate Ligament Reconstruction: A Prospective, Single-Blinded, Randomized Trial. *Orthop J Sport*

- Med. 2018 Sep 1;6(9).
179. Ghodki P, Shalu P, Sardesai S. Ultrasound-guided adductor canal block versus femoral nerve block for arthroscopic anterior cruciate ligament repair under general anesthesia. *J Anaesthesiol Clin Pharmacol*. 2018 Apr 1;34(2):242–6.
  180. Stebler K, Martin R, Kirkham KR, Lambert J, De Sede A, Albrecht E. Adductor canal block versus local infiltration analgesia for postoperative pain after anterior cruciate ligament reconstruction: a single centre randomised controlled triple-blinded trial. *Br J Anaesth*. 2019 Aug 1;123(2):e343–9.
  181. Kejriwal R, Cooper J, Legg A, Stanley J, Rosenfeldt MP, Walsh SJ. Efficacy of the Adductor Canal Approach to Saphenous Nerve Block for Anterior Cruciate Ligament Reconstruction With Hamstring Autograft: A Randomized Controlled Trial. *Orthop J Sport Med*. 2018 Oct 1;6(10).
  182. Sehmbi H, Brull R, Shah UJ, El-Boghdadly K, Nguyen D, Joshi GP, et al. Evidence Basis for Regional Anesthesia in Ambulatory Arthroscopic Knee Surgery and Anterior Cruciate Ligament Reconstruction: Part II: Adductor Canal Nerve Block-A Systematic Review and Meta-analysis. *Anesth Analg*. 2019 Feb 1;128(2):223–38.
  183. Yung EM, Brull R, Albrecht E, Joshi GP, Abdallah FW. Evidence basis for regional anesthesia in ambulatory anterior cruciate ligament reconstruction: Part III: Local instillation analgesia-A systematic review and meta-Analysis. Vol. 128, *Anesthesia and Analgesia*. Lippincott Williams and Wilkins; 2019. p. 426–37.
  184. McCartney CJL, Mclsaac D. Are Peripheral Nerve Blocks Indicated in Ambulatory Knee Surgery? *Anesth Analg* [Internet]. 2019 Jan [cited 2020 Jan 11];128(1):3–4. Available from: <http://insights.ovid.com/crossref?an=00000539-201901000-00003>
  185. Piper SL, Kim HT. Comparison of ropivacaine and bupivacaine toxicity in human articular chondrocytes. *J Bone Jt Surg - Ser A*. 2008;90(5):986–91.
  186. Breu A, Rosenmeier K, Kujat R, Angele P, Zink W. The cytotoxicity of bupivacaine, ropivacaine, and mepivacaine on human chondrocytes and cartilage. *Anesth Analg*. 2013 Aug;117(2):514–22.
  187. Jansson H, Narvy SJ, Mehran N. Perioperative pain management strategies for anterior cruciate ligament reconstruction. *JBJS Rev*. 2018 Mar 1;6(3).
  188. Baverel L, Cucurulo T, Lutz C, Colombet, Cournapeau J, Dalmay F, et al. Anesthesia and

- analgesia methods for outpatient anterior cruciate ligament reconstruction. *Orthop Traumatol Surg Res*. 2016 Dec 1;102(8):S251–5.
189. Kraus Schmitz J, Lindgren V, Janarv PM, Forssblad M, St Iman A. Deep venous thrombosis and pulmonary embolism after anterior cruciate ligament reconstruction. *Bone Jt J*. 2019 Jan 1;101B(1):34–40.
  190. Erickson BJ, Saltzman BM, Campbell KA, Fillingham YA, Harris JD, Gupta AK, et al. Rates of Deep Venous Thrombosis and Pulmonary Embolus After Anterior Cruciate Ligament Reconstruction: A Systematic Review. *Sports Health [Internet]*. 2015 May [cited 2020 Jan 12];7(3):261–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26131305>
  191. Dumville JC, Mcfarlane E, Edwards P, Lipp A, Holmes A, Liu Z. Preoperative skin antiseptics for preventing surgical wound infections after clean surgery. Vol. 2017, *Cochrane Database of Systematic Reviews*. John Wiley and Sons Ltd; 2015.
  192. Samuelsen BT, Webster KE, Johnson NR, Hewett TE, Krych AJ. Hamstring Autograft versus Patellar Tendon Autograft for ACL Reconstruction: Is There a Difference in Graft Failure Rate? A Meta-analysis of 47,613 Patients. *Clin Orthop Relat Res [Internet]*. 2017 Oct [cited 2020 Jan 12];475(10):2459–68. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28205075>
  193. Slone HS, Romine SE, Premkumar A, Xerogeanes JW. Quadriceps tendon autograft for anterior cruciate ligament reconstruction: a comprehensive review of current literature and systematic review of clinical results. *Arthroscopy [Internet]*. 2015 Mar [cited 2020 Jan 12];31(3):541–54. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25543249>
  194. Hardy A, Casabianca L, Andrieu K, Baverel L, Noailles T, Junior French Arthroscopy Society. Complications following harvesting of patellar tendon or hamstring tendon grafts for anterior cruciate ligament reconstruction: Systematic review of literature. *Orthop Traumatol Surg Res [Internet]*. 2017 [cited 2020 Jan 12];103(8S):S245–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28888527>
  195. Xie X, Liu X, Chen Z, Yu Y, Peng S, Li Q. A meta-analysis of bone-patellar tendon-bone autograft versus four-strand hamstring tendon autograft for anterior cruciate ligament reconstruction. Vol. 22, *Knee*. Elsevier; 2015. p. 100–10.
  196. Widner M, Dunleavy M, Lynch S. Outcomes Following ACL Reconstruction Based on Graft Type: Are all Grafts Equivalent? *Curr Rev Musculoskelet Med [Internet]*. 2019 Nov 16 [cited 2020 Jan 12]; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/31734844>

197. Cavaignac E, Coulin B, Tscholl P, Nik Mohd Fatmy N, Duthon V, Menetrey J. Is Quadriceps Tendon Autograft a Better Choice Than Hamstring Autograft for Anterior Cruciate Ligament Reconstruction? *Am J Sports Med* [Internet]. 2017 [cited 2020 Jan 12];036354651668866. Available from: <http://journals.sagepub.com/doi/10.1177/0363546516688665>
198. Hurley ET, Calvo-Gurry M, Withers D, Farrington SK, Moran R, Moran CJ. Quadriceps Tendon Autograft in Anterior Cruciate Ligament Reconstruction: A Systematic Review. *Arthroscopy* [Internet]. 2018 [cited 2020 Jan 12];34(5):1690–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29628380>
199. Lind M, Strauss MJ, Nielsen T, Engebretsen L. Quadriceps tendon autograft for anterior cruciate ligament reconstruction is associated with high revision rates: results from the Danish Knee Ligament Registry. *Knee Surg Sports Traumatol Arthrosc* [Internet]. 2019 Oct 22 [cited 2020 Jan 12]; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/31641810>
200. Tejwani SG, Chen J, Funahashi TT, Love R, Maletis GB. Revision risk after allograft anterior cruciate ligament reconstruction: Association with graft processing techniques, patient characteristics, and graft type. *Am J Sport Med* [Internet]. 2015 [cited 2020 Jan 12];43(11):2696–705. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26068037>
201. Brown MJ, Carter T. ACL allograft: Advantages and when to use. *Sports Med Arthrosc*. 2018;26(2):75–8.
202. Kaeding CC, Pedroza AD, Reinke EK, Huston LJ, Hewett TE, Flanigan DC, et al. Change in Anterior Cruciate Ligament Graft Choice and Outcomes Over Time. *Arthrosc - J Arthrosc Relat Surg*. 2017 Nov 1;33(11):2007–14.
203. Faunø P, Rahr-Wagner L, Lind M. Risk for revision after anterior cruciate ligament reconstruction is higher among adolescents: Results from the Danish registry of knee ligament reconstruction. *Orthop J Sport Med*. 2014;2(10):1–7.
204. Peltola EK, Lindahl J, Koskinen SK. The reverse Segond fracture: not associated with knee dislocation and rarely with posterior cruciate ligament tear. *Emerg Radiol* [Internet]. 2014 Jun 7 [cited 2018 May 28];21(3):245–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24395399>
205. Offerhaus C, Balke M, Hente J, Gehling M, Blendl S, Höher J. Vancomycin pre-soaking of the graft reduces postoperative infection rate without increasing risk of graft failure and arthrofibrosis in ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* [Internet]. 2019 Sep

- [cited 2020 Jan 17];27(9):3014–21. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30666370>
206. Naendrup J-H, Marche B, de Sa D, Koenen P, Otchwemah R, Wafaisade A, et al. Vancomycin-soaking of the graft reduces the incidence of septic arthritis following ACL reconstruction: results of a systematic review and meta-analysis. *Knee Surg Sports Traumatol Arthrosc* [Internet]. 2019 Jan 17 [cited 2020 Jan 17]; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30656372>
207. Phegan M, Grayson JE, Vertullo CJ. No infections in 1300 anterior cruciate ligament reconstructions with vancomycin pre-soaking of hamstring grafts. *Knee Surg Sports Traumatol Arthrosc* [Internet]. 2016 Sep [cited 2020 Jan 17];24(9):2729–35. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25771788>
208. Figueroa D, Figueroa F, Calvo R, Lopez M, Goñi I. Presoaking of Hamstring Autografts in Vancomycin Decreases the Occurrence of Infection Following Primary Anterior Cruciate Ligament Reconstruction. *Orthop J Sport Med* [Internet]. 2019 Sep [cited 2020 Jan 17];7(9):2325967119871038. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/31598528>
209. Schuster P, Schlumberger M, Mayer P, Eichinger M, Geßlein M, Richter J. Soaking of autografts in vancomycin is highly effective in preventing postoperative septic arthritis after revision anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* [Internet]. 2019 Dec 3 [cited 2020 Jan 17]; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/31797021>
210. Pérez-Prieto D, Torres-Claramunt R, Gelber PE, Shehata TMA, Pelfort X, Monllau JC. Autograft soaking in vancomycin reduces the risk of infection after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* [Internet]. 2016 Sep [cited 2020 Jan 17];24(9):2724–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25416672>
211. Baron JE, Shamrock AG, Cates WT, Cates RA, An Q, Wolf BR, et al. Graft Preparation with Intraoperative Vancomycin Decreases Infection After ACL Reconstruction. *J Bone Jt Surg*. 2019 Oct;101(24):2187–93.
212. Schüttler K-F, Scharm A, Stein T, Heyse TJ, Lohoff M, Sommer F, et al. Biomechanical and microbiological effects of local vancomycin in anterior cruciate ligament (ACL) reconstruction: a porcine tendon model. *Arch Orthop Trauma Surg* [Internet]. 2019 Jan [cited 2020 Jan

- 17];139(1):73–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30039308>
213. Grayson JE, Grant GD, Dukie S, Vertullo CJ. The in vitro elution characteristics of vancomycin from tendons. *Clin Orthop Relat Res*. 2011;469(10):2948–52.
214. Pérez-Prieto D, Portillo ME, Torres-Claramunt R, Pelfort X, Hinarejos P, Monllau JC. Contamination occurs during ACL graft harvesting and manipulation, but it can be easily eradicated. *Knee Surg Sports Traumatol Arthrosc* [Internet]. 2018 Feb [cited 2020 Jan 17];26(2):558–62. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28988312>
215. Jefferies JG, Aithie JMS, Spencer SJ. Vancomycin-soaked wrapping of harvested hamstring tendons during anterior cruciate ligament reconstruction. A review of the ‘vancomycin wrap.’ *Knee*. 2019. p. 26 (3):524–529.
216. Wilson WT, Hopper GP, Byrne PA, MacKay GM. Anterior Cruciate Ligament Repair with Internal Brace Ligament Augmentation. *Surg Technol Int*. 2016 Oct 26;29:273–8.
217. Sonnery-Cottet B, Colombet P. Partial tears of the anterior cruciate ligament. Vol. 102, *Orthopaedics and Traumatology: Surgery and Research*. Elsevier Masson SAS; 2016. p. S59–67.
218. Papalia R, Franceschi F, Zampogna B, Tecame A, Maffulli N, Denaro V. Surgical management of partial tears of the anterior cruciate ligament. Vol. 22, *Knee Surgery, Sports Traumatology, Arthroscopy*. 2014. p. 154–65.
219. Murray MM, Kalish LA, Fleming BC, Flutie B, Freiburger C, Henderson RN, et al. Bridge-Enhanced Anterior Cruciate Ligament Repair: Two-Year Results of a First-in-Human Study. *Orthop J Sport Med*. 2019 Mar 1;7(3).
220. van Eck CF, Limpisvasti O, ElAttrache NS. Is There a Role for Internal Bracing and Repair of the Anterior Cruciate Ligament? A Systematic Literature Review. Vol. 46, *American Journal of Sports Medicine*. SAGE Publications Inc.; 2018. p. 2291–8.
221. van der List JP, DiFelice GS. Primary repair of the anterior cruciate ligament: A paradigm shift. Vol. 15, *Surgeon*. Elsevier Ltd; 2017. p. 161–8.
222. Riediger MD, Stride D, Coke SE, Kurz AZ, Duong A, Ayeni OR. ACL Reconstruction with Augmentation: a Scoping Review. Vol. 12, *Current Reviews in Musculoskeletal Medicine*. Humana Press Inc.; 2019. p. 166–72.
223. Paschos NK, Howell SM. Anterior cruciate ligament reconstruction: Principles of treatment.

- EFORT Open Rev. 2016 Nov 1;1(11):398–408.
224. Kraeutler MJ, Patel K V, Hosseini A, Li G, Gill TJ, Bravman JT. Variability in the Clock Face View Description of Femoral Tunnel Placement in ACL Reconstruction Using MRI-Based Bony Models. *J Knee Surg* [Internet]. 2018 Nov [cited 2020 Jan 13];31(10):965–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29421839>
  225. McConkey MO, Amendola A, Ramme AJ, Dunn WR, Flanigan DC, Britton CL, et al. Arthroscopic agreement among surgeons on anterior cruciate ligament tunnel placement. *Am J Sports Med* [Internet]. 2012 Dec [cited 2020 Jan 13];40(12):2737–46. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23075806>
  226. Han Y, Hart A, Martineau PA. Is the clock face an accurate, precise, and reliable measuring tool for anterior cruciate ligament reconstruction? *Arthroscopy* [Internet]. 2014 Jul [cited 2020 Jan 13];30(7):849–55. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24821225>
  227. Dimitriou D, Wang Z, Zou D, Tsai T-Y, Helmy N. The Femoral Footprint Position of the Anterior Cruciate Ligament Might Be a Predisposing Factor to a Noncontact Anterior Cruciate Ligament Rupture. *Am J Sports Med* [Internet]. 2019 Dec [cited 2020 Jan 13];47(14):3365–72. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/31647682>
  228. Parkinson B, Gogna R, Robb C, Thompson P, Spalding T. Anatomic ACL reconstruction: the normal central tibial footprint position and a standardised technique for measuring tibial tunnel location on 3D CT. *Knee Surgery, Sport Traumatol Arthrosc*. 2017 May 1;25(5):1568–75.
  229. Eysturoy NH, Nielsen TG, Lind MC. Anteromedial Portal Drilling Yielded Better Survivorship of Anterior Cruciate Ligament Reconstructions When Comparing Recent Versus Early Surgeries With This Technique. *Arthrosc - J Arthrosc Relat Surg*. 2019 Jan 1;35(1):182–9.
  230. Arno S, Bell CP, Alaia MJ, Singh BC, Jazrawi LM, Walker PS, et al. Does Anteromedial Portal Drilling Improve Footprint Placement in Anterior Cruciate Ligament Reconstruction? *Clin Orthop Relat Res*. 2016 Jul 1;474(7):1679–89.
  231. Eysturoy NH, Nissen KA, Nielsen T, Lind M. The Influence of Graft Fixation Methods on Revision Rates After Primary Anterior Cruciate Ligament Reconstruction. *Am J Sports Med*. 2018 Mar 1;46(3):524–30.
  232. Debieux P, Franciozi CE, Lenza M, Tamaoki MJ, Magnussen RA, Faloppa F, et al. Bioabsorbable versus metallic interference screws for graft fixation in anterior cruciate ligament

- reconstruction. *Cochrane Database Syst Rev* [Internet]. 2016 Jul 24 [cited 2018 Nov 12];7:CD009772. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27450741>
233. Sonnery-Cottet B, Daggett M, Fayard J-M, Ferretti A, Helito CP, Lind M, et al. Anterolateral Ligament Expert Group consensus paper on the management of internal rotation and instability of the anterior cruciate ligament - deficient knee. *J Orthop Traumatol* [Internet]. 2017 Jun 20 [cited 2019 Sep 21];18(2):91–106. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28220268>
234. Getgood A, Brown C, Lording T, Amis A, Claes S, Geeslin A, et al. The anterolateral complex of the knee: results from the International ALC Consensus Group Meeting. *Knee Surgery, Sport Traumatol Arthrosc*. 2019 Jan 30;27(1):166–76.
235. Lutz C. Role of anterolateral reconstruction in patients undergoing anterior cruciate ligament reconstruction. *Orthop Traumatol Surg Res* [Internet]. 2018 Feb [cited 2019 Sep 21];104(1):S47–53. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29191469>
236. Herbst E, Arilla F V., Guenther D, Yacuzzi C, Rahnama-Azar AA, Fu FH, et al. Lateral Extra-articular Tenodesis Has No Effect in Knees With Isolated Anterior Cruciate Ligament Injury. *Arthrosc J Arthrosc Relat Surg* [Internet]. 2018 Jan [cited 2019 Sep 21];34(1):251–60. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29079261>
237. Geeslin AG, Moatshe G, Chahla J, Kruckeberg BM, Muckenhirn KJ, Dornan GJ, et al. Anterolateral Knee Extra-articular Stabilizers: A Robotic Study Comparing Anterolateral Ligament Reconstruction and Modified Lemaire Lateral Extra-articular Tenodesis. *Am J Sports Med* [Internet]. 2018 Mar 21 [cited 2019 Sep 21];46(3):607–16. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29268024>
238. Claes S, Vereecke E, Maes M, Victor J, Verdonk P, Bellemans J. Anatomy of the anterolateral ligament of the knee. *J Anat* [Internet]. 2013 Oct [cited 2018 Jul 7];223(4):321–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23906341>
239. Musahl V, Herbst E, Burnham JM, Fu FH. The Anterolateral Complex and Anterolateral Ligament of the Knee. *J Am Acad Orthop Surg* [Internet]. 2018 Apr 15 [cited 2018 Jul 7];26(8):261–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29517517>
240. Lowe WR, Warth RJ, Davis EP, Bailey L. Functional Bracing After Anterior Cruciate Ligament Reconstruction: A Systematic Review. *J Am Acad Orthop Surg* [Internet]. 2017 Mar [cited 2020 Jan 5];25(3):239–49. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28195986>



241. Rodríguez-Merchán EC. Knee Bracing After Anterior Cruciate Ligament Reconstruction. *Orthopedics* [Internet]. 2016 Jul 1 [cited 2020 Jan 12];39(4):e602-9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27203412>
242. Bordes P, Laboute E, Bertolotti A, Dalmay JF, Puig P, Trouve P, et al. No beneficial effect of bracing after anterior cruciate ligament reconstruction in a cohort of 969 athletes followed in rehabilitation. *Ann Phys Rehabil Med*. 2017 Jul 1;60(4):230–6.
243. Grant JA. Updating Recommendations for Rehabilitation after ACL Reconstruction: a Review. *Clin J Sport Med* [Internet]. 2013 Nov [cited 2020 Jan 12];23(6):501–2. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24169301>
244. Andrade R, Pereira R, van Cingel R, Staal JB, Espregueira-Mendes J. How should clinicians rehabilitate patients after ACL reconstruction? A systematic review of clinical practice guidelines (CPGs) with a focus on quality appraisal (AGREE II). *Br J Sports Med* [Internet]. 2019 Jun 7 [cited 2020 Jan 12]; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/31175108>
245. Wright RW, Fetzer GB. Bracing after ACL reconstruction: A systematic review. In: *Clinical Orthopaedics and Related Research*. Lippincott Williams and Wilkins; 2007. p. 162–8.
246. Wright RW. Rehabilitation predictors of clinical outcome following revision ACL reconstruction in the MARS cohort. *J Bone Jt Surg - Am Vol*. 2019 May 1;101(9):779–86.
247. Herrington L, Myer G HI. Task based rehabilitation protocol for elite athletes following Anterior Cruciate ligament reconstruction: a clinical commentary. *Phys Ther Sport*. 2013;14:188–98.
248. Adams D, Logerstedt D, Hunter-Giordano A, Axe MJ, Snyder-Mackler L. Current concepts for anterior cruciate ligament reconstruction: A criterion-based rehabilitation progression. *J Orthop Sports Phys Ther*. 2012;42(7):601–14.
249. Giotis D, Paschos NK, Zampeli F, Pappas E, Mitsionis G, Georgoulis AD. Bracing can partially limit tibial rotation during stressful activities after anterior crucial ligament reconstruction with a hamstring graft. *Orthop Traumatol Surg Res*. 2016 Sep 1;102(5):601–6.
250. Perrone GS, Webster KE, Imbriaco C, Portilla GM, Vairagade A, Murray MM, et al. Risk of Secondary ACL Injury in Adolescents Prescribed Functional Bracing After ACL Reconstruction. *Orthop J Sport Med*. 2019;7(11).
251. (No Title) [Internet]. [cited 2020 Jan 12]. Available from: <https://www.aclregister.nu/media/uploads/Annual>

reports/annual\_report\_swedish\_acl\_registry\_2018.pdf

252. (No Title) [Internet]. [cited 2020 Jan 12]. Available from: [https://reports.njrcentre.org.uk/Portals/0/PDFdownloads/NJR 16th Annual Report 2019.pdf](https://reports.njrcentre.org.uk/Portals/0/PDFdownloads/NJR%2016th%20Annual%20Report%202019.pdf)
253. de Jong S, van Caspel D, van Haeff M SD. Functional assessment and muscle strength before and after reconstruction of chronic anterior cruciate ligament lesions. *Arthroscopy*. 2007;23:21–28.
254. Grindem H, Granan LP, Risberg MA, Engebretsen L, Snyder-Mackler L, Eitzen I. How does a combined preoperative and postoperative rehabilitation programme influence the outcome of ACL reconstruction 2 years after surgery? A comparison between patients in the Delaware-Oslo ACL Cohort and the Norwegian National Knee Ligament Registry. *Br J Sports Med*. 2015;49(6):385–389.
255. Lynch A, Logerstedt D, Axe M S-ML. Quadriceps activation failure after anterior cruciate ligament rupture is not mediated by knee joint effusion. *J Orthop Sports Phys Ther*. 2012;42:502–10.
256. Decker M, Torry M, Noonan T, Sterett W SJ. Gait retraining after anterior cruciate ligament reconstruction. *Arch Phys Med Rehabil*. 2004;85:848–56.
257. Myer G, Brent J, Ford K & HT. A pilot study to determine the effect of trunk and hip focused neuromuscular training on hip and knee isokinetic strength. *Br J Sports Med*. 2008;42:614–9.
258. Paterno M, Schmitt L, Ford K, Rauh M, Myer G, Huang B HT. Biomechanical measures during landing and postural stability predict second anterior cruciate ligament injury after anterior cruciate reconstruction and return to sport. *Am J Sports Med*. 2010;38:1968–78.
259. Angelozzi M, Madama M, Corsica C, Calvisi V, Properzi G, McCaw ST, et al. Rate of force development as an adjunctive outcome measure for return-to-sport decisions after anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther* [Internet]. 2012 Sep [cited 2019 Dec 31];42(9):772–80. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22814219>
260. Myer GD, Martin L, Ford KR, Paterno M V., Schmitt LC, Heidt RS, et al. No association of time from surgery with functional deficits in athletes after anterior cruciate ligament reconstruction: Evidence for objective return-to-sport criteria. *Am J Sports Med*. 2012 Oct;40(10):2256–63.
261. Trulsson A, Roos E, Ageberg E GM, Trulsson A, Roos EM, Ageberg E, Garwicz M. Relationship between postural orientation and self-reported function, hop performance and muscle power

- in subjects with anterior cruciate ligament injury. *BMC Musculoskelet Disord.* 2010;11:143–15.
262. Reinke E, Spindler K, Lorrington D, Jones M, Schmitz L, Flanigan D, Qi A, Quiram A, Preston E, Martin M, Schroeder B, Parker R, Kaeding C, Borzi L, Pedroza A, Huston L, Harrell F DW. Hop tests correlate with IKDC and KOOS at minimum of 2 years after primary ACL reconstructions. *Knee Surg Sport Traumatol Arthrosc.* 2011;19:1806–16.
263. Barber-Westin SD, Noyes FR. Factors used to determine return to unrestricted sports activities after anterior cruciate ligament reconstruction. *Arthrosc - J Arthrosc Relat Surg.* 2011;27(12):1697–705.
264. Grindem H, Snyder-Mackler L, Moksnes H, Engebretsen L, Risberg MA. Simple decision rules can reduce reinjury risk by 84% after ACL reconstruction: The Delaware-Oslo ACL cohort study. *Br J Sports Med.* 2016 Jul 1;50(13):804–8.
265. Herrington L, Myer G, Horsley I. Task based rehabilitation protocol for elite athletes following Anterior Cruciate ligament reconstruction: A clinical commentary. Vol. 14, *Physical Therapy in Sport.* 2013. p. 188–98.
266. Ardern CL, Glasgow P, Schneiders A, Witvrouw E, Clarsen B, Cools A, et al. 2016 Consensus statement on return to sport from the First World Congress in Sports Physical Therapy, Bern. *Br J Sports Med.* 2016 Jul 1;50(14):853–64.
267. Webster KE, Feller JA, Lambros C. Development and preliminary validation of a scale to measure the psychological impact of returning to sport following anterior cruciate ligament reconstruction surgery. *Phys Ther Sport.* 2008 Feb;9(1):9–15.